

Homogeneity of proper complex equifocal submanifolds

Naoyuki Koike

*Department of Mathematics, Faculty of Science, Tokyo University of Science,
26 Wakamiya-cho Shinjuku-ku, Tokyo 162-8601 Japan
E-mail address: koike@ma.kagu.tus.ac.jp*

Abstract

In a symmetric space of non-compact type, the notion of a complex equifocal submanifold (with flat section) is defined. It is conjectured that this notion coincides with the notion of an isoparametric submanifold with flat section. As a subclass of a complex equifocal submanifold, the notion of a proper complex equifocal submanifold is defined. In this paper, we show that all irreducible proper complex equifocal submanifolds of codimension greater than one in a symmetric space of non-compact type are extrinsically homogeneous and hence they occur as principal orbits of complex hyperpolar actions on the symmetric space. The proof is performed by showing the homogeneity of the lifted submanifold of the complexification of the original submanifold to an infinite dimensional anti-Kaehlerian space and inducing the homogeneity of the original submanifold from the homogeneity of the lifted submanifold.

Keywords: proper complex equifocal submanifold, proper anti-Kaehlerian isoparametric submanifold, complex principal curvature, complex curvature distribution

1 Introduction

C.L. Terng and G. Thorbergsson [TT] introduced the notion of an equifocal submanifold in a Riemannian symmetric space, which is defined as a compact submanifold with globally flat and abelian normal bundle such that the focal radii for each parallel normal vector field are constant. This notion is a generalization of isoparametric submanifolds in the Euclidean space and isoparametric hypersurfaces in the sphere or the hyperbolic space. For (not necessarily compact) submanifolds in a Riemannian symmetric space of non-compact type, the equifocality is a rather weak property. So, we [Koi1,2] introduced the

notion of a complex focal radius as a general notion of a focal radius and defined the notion of a complex equifocal submanifold as a submanifold with globally flat and abelian normal bundle such that the complex focal radii for each parallel normal vector field are constant and that they have constant multiplicities. Here we note that the notion of a complex focal radius (hence the notion of the complex equifocality) can be defined for submanifolds in a general symmetric space but, in the case where the ambient symmetric space is of non-negative curvature, all complex focal radii are real, that is, they are focal radii and hence the complex equifocality is equivalent to the equifocality. E. Heintze, X. Liu and C. Olmos [HLO] defined the notion of an isoparametric submanifold with flat section as a submanifold with globally flat and abelian normal bundle such that the sufficiently close parallel submanifolds are of constant mean curvature with respect to the radial direction. The following fact is known (see Theorem 15 of [Koi2]):

All isoparametric submanifolds with flat section in a symmetric space of non-compact type are complex equifocal and, conversely, all curvature-adapted and complex equifocal submanifolds are isoparametric submanifolds with flat section.

Here the curvature-adaptedness means that, for any normal vector v , $R(\cdot, v)v$ preserves the tangent space invariantly and it commutes with the shape operator A_v . Furthermore, as its subclass, we [Koi1,2] introduced the notion of a proper complex equifocal submanifold as a complex equifocal submanifold whose complex focal structure has a certain kind of regularity. Here we note that the notion of a proper complex equifocal submanifold in a general symmetric space can be defined but, in the case where the ambient symmetric space is of compact type, it is easy to show that this notion coincides with the notion of a complex equifocal submanifold. In [Koi8], we showed that, for a curvature-adapted complex equifocal submanifold M , it is proper complex equifocal if and only if it admits no non-Euclidean type focal point on the ideal boundary of the ambient symmetric space, where the notion of a non-Euclidean type focal point on the ideal boundary was introduced in [Koi8]. Therefore, for a submanifold M in a symmetric space of non-compact type, the following two statements are equivalent:

- (I) M is curvature-adapted and proper complex equifocal.
- (II) M is a curvature-adapted isoparametric submanifold with flat section admitting no non-Euclidean type focal point on the ideal boundary.

Let G/K be a symmetric space of non-compact type and H be a symmetric subgroup of G (i.e., there exists an involution σ of G with $(\text{Fix } \sigma)_0 \subset H \subset \text{Fix } \sigma$, where $\text{Fix } \sigma$ is the fixed point group of σ and $(\text{Fix } \sigma)_0$ is the identity component of $\text{Fix } \sigma$). Then the H -action on G/K is called a *Hermann type action*. For this action, the following fact is known ([Koi3]):

Principal orbits of Hermann type actions are curvature-adapted and proper complex equifocal.

For a (general) submanifold in a Riemannian symmetric space of non-compact type, the (non-real) complex focal radii are defined algebraically. We needed to find their geometrical essence. For its purpose, we defined the complexification of the ambient Riemannian symmetric space and the extrinsic complexification of the submanifold as a certain kind of submanifold in the complexified symmetric space, where the original submanifold needs to be assumed to be complete and real analytic. In the sequel, we assume that all submanifolds in the Riemannian symmetric space are complete and real analytic. We [Koi2] showed that the complex focal radii of the original submanifold indicate the positions of the focal points of the complexified submanifold. If the original submanifold is complex equifocal, then the extrinsic complexification is an anti-Kaehlerian equifocal submanifold in the sense of [Koi2]. Also, if the original submanifold is proper complex equifocal, then the complexified one is a proper anti-Kaehlerian equifocal submanifold in the sense of this paper. Thus, the study of an anti-Kaehlerian equifocal (resp. proper anti-Kaehlerian equifocal) submanifold leads to that of a complex equifocal (resp. proper complex equifocal) submanifold. The complexified submanifold is not necessarily complete. In the global research, we need to extend the complexified submanifold to a complete one. In [Koi2] and [Koi6], we constructed the complete extension of the complexified submanifold in mutually different methods. In 1999, E. Heintze and X. Liu [HL2] showed that all irreducible isoparametric submanifolds of codimension greater than one in the (separable) Hilbert space are extrinsically homogeneous, which is the infinite dimensional version of the homogeneity theorem for isoparametric submanifolds in a (finite dimensional) Euclidean space by G. Thorbergsson. Note that the result of Thorbergsson states that all irreducible isoparametric submanifolds of codimension greater than two in a Euclidean space are extrinsically homogeneous. In 2002, by using the result of Heintze-Liu, U. Christ [Ch] showed that all irreducible equifocal submanifolds of codimension greater than one in a simply connected symmetric space of compact type are extrinsically homogeneous. The outline of the proof is as follows. For an irreducible equifocal submanifold M of codimension greater than one in a simply connected symmetric space G/K of compact type, any component $\widetilde{M}$ of the inverse image $(\pi \circ \phi)^{-1}(M)$ of M by the composition of the parallel transport map $\phi : H^0([0, 1], \mathfrak{g}) \rightarrow G$ with the natural projection $\pi : G \rightarrow G/K$ is an irreducible isoparametric submanifold of codimension greater than one in the Hilbert space $H^0([0, 1], \mathfrak{g})$, where $H^0([0, 1], \mathfrak{g})$ is the space of all L^2 -integrable paths in $\mathfrak{g} := \text{Lie } G$. By the result of Heintze-Liu, $\widetilde{M}$ is extrinsically homogeneous, that is, $\widetilde{M}$ is an orbit of some subgroup H of the isometry group of $H^0([0, 1], \mathfrak{g})$. From the H -action on $H^0([0, 1], \mathfrak{g})$, he constructed a subgroup $\overline{H}$ of G such that M is an orbit of the $\overline{H}$ -action on G/K . Hence M is extrinsically homogeneous.

For a complex equifocal submanifold M in a symmetric space G/K of non-compact type, the complete extrinsically complexification M^c of M is defined as an anti-Kaehlerian submanifold in the anti-Kaehlerian symmetric space G^c/K^c . Without loss of generality,

we may assume that K (hence K^c) is connected and that G^c is simply connected. Let π^c be the natural projection of G^c onto G^c/K^c and $\phi^c : H^0([0, 1], \mathfrak{g}^c) \rightarrow G^c$ be the parallel transport map for G^c . Then $\widetilde{M}^c := (\pi^c \circ \phi^c)^{-1}(M^c)$ is an anti-Kaehlerian isoparametric submanifold in the infinite dimensional anti-Kaehlerian space $H^0([0, 1], \mathfrak{g}^c)$ in the sense of [Koi2]. In particular, if M is proper complex equifocal, then $\widetilde{M}^c$ is a proper anti-Kaehlerian isoparametric submanifold in the sense of [Koi2]. First we prove the following fact for the homogeneity of the lifted submanifold $\widetilde{M}^c$:

If M is an irreducible proper complex equifocal submanifold of codimension greater than one in a symmetric space of non-compact type, then the lifted submanifold $\widetilde{M}^c$ is extrinsically homogeneous.

By using this fact, we prove the following homogeneity theorem for an irreducible proper complex equifocal submanifold.

Theorem A. *All irreducible proper complex equifocal submanifolds of codimension greater than one in a symmetric space of non-compact type are extrinsically homogeneous.*

Let G/K be a symmetric space of non-compact type and H be a closed subgroup of G which admits an embedded flat submanifold of G/K meeting all H -orbits orthogonally. Then the H -action is called a *complex hyperpolar action*. From Theorem A, we can show the following fact.

Corollary B. *Let M be an irreducible proper complex equifocal submanifold of codimension greater than one in a symmetric space G/K of non-compact type. Then the following statements (i) and (ii) hold:*

- (i) *M occurs as a principal orbit of a complex hyperpolar action on G/K .*
- (ii) *If M admits a totally geodesic focal submanifold, then it occurs as a principal orbit of a Hermann type action on G/K .*

The outline of the discussions in Sections 3 and 4 In Section 3, for a submanifold M as in the statement of Theorem A, we prove that $\widetilde{M}^c := (\pi^c \circ \phi^c)^{-1}(M^c)$ is homogeneous by imitating the proof of the homogeneity of an irreducible infinite dimensional isoparametric submanifold of codimension greater than one in a Hilbert space by Heintze-Liu [HL2]. In the proof of the homogeneity of the isoparametric submanifold, Heintze-Liu [HL2] used the curvature distributions on the isoparametric submanifold. On the other hand, $\widetilde{M}^c$ is proper anti-Kaehlerian isoparametric submanifold and hence the complex curvature distributions on $\widetilde{M}^c$ is defined. See the next section about the definition of the complex curvature distribution. In the proof of the homogeneity of $\widetilde{M}^c$, we use the complex curvature distributions. We sketch the outline of the discussion in Section 3. First we recall the generalized Chow's theorem for the set of reachable points for a certain kind of

family of local vector fields on a Hilbert manifold, which were proved in [HL2]. By using this theorem, we show that the set (which is denoted by $Q(u_0)$) of all the points connected with a fixed point u_0 of $\widetilde{M}^c$ by piecewise smooth curves each of whose segment is contained in a complex curvature spheres (the integral manifolds of the complex curvature distributions) is dense in $\widetilde{M}^c$ (see Proposition 3.2). Next we show the homogeneous slice theorem (Theorem 3.3) for $\widetilde{M}^c$ by using the homogeneous slice theorem for M^c , which was proved in [Koi6]. Next we construct a holomorphic isometry F_γ of $V := H^0([0, 1], \mathfrak{g}^c)$ with $F_\gamma(\gamma(0)) = \gamma(1)$ for a certain kind of piecewise smooth curve $\gamma : [0, 1] \rightarrow \widetilde{M}^c$ (which is a special one of curves called a E_i -horizontal curve) in $\widetilde{M}^c$, where we note that E_i is a complex curvature distribution. This holomorphic isometry F_γ is constructed by using the fact that, for each linear isometry of the tangent space at a point u_0 of V onto the tangent space at another point u_1 of V , there (uniquely) exists a holomorphic isometry of V whose differential at u_0 coincides with the linear isometry (see (3.1)). Here we note that this fact holds because V is an (anti-Kaehlerian) linear space. In the discussion in Section 3, it is a key to show that F_γ preserves $\widetilde{M}^c$ invariantly (see Proposition 3.5). In this proof, we use the assumptions of the irreducibility of M and $\text{codim } M \geq 2$, and the above homogeneous slice theorem for $\widetilde{M}^c$ (see the proof of Lemma 3.5.1). Next we show that, for each $u \in Q(u_0)$, there exists a holomorphic isometry f of V with $f(u_0) = u$ and $f(\widetilde{M}^c) = \widetilde{M}^c$ (see Proposition 3.6). Finally, by using Propositions 3.2 and 3.6, we show that $\widetilde{M}^c$ is homogeneous. In Section 4, we induce the homogeneity of M from the homogeneity of $\widetilde{M}^c$ by imitating the proof of the homogeneity of an irreducible equifocal submanifold of codimension greater than one in a symmetric space of compact type by Christ [Ch]. Here we sketch the outline of the discussion in Section 4. The group $H^1([0, 1], G^c)$ acts on V as the gauge action. Denote by $\rho : H^1([0, 1], G^c) \rightarrow I_h(V)$ the representation on V associated with this action, where $I_h(V)$ is the holomorphic isometry group of V . In the discussion in Section 4, it is a key to show that the group $H' := \{f \in I_h(V) \mid f(\widetilde{M}^c) = \widetilde{M}^c\}$ is contained in $\rho(H^1([0, 1], G^c))$. This fact is shown by comparing the spaces of Killing fields corresponding to H' and $\rho(H^1([0, 1], G^c))$ (see the proofs of Lemmas 4.1.1 $\sim$ 4.1.11). Thus the group H' is given as the image $\rho(H)$ of some subgroup H of $H^1([0, 1], G^c)$. Since $\widetilde{M}^c$ is homogeneous, we have $\widetilde{M}^c = \rho(H) \cdot u_0$ ($u_0 \in \widetilde{M}^c$). We construct a subgroup $\overline{H}$ of $G^c \times G^c$ from H and show $\overline{H} \cdot g_0 = \pi^{c-1}(M^c)$ ($g_0 \in \pi^{c-1}(M^c)$). Next we construct a subgroup $\overline{H}_{\mathbf{R}}$ of $G \times G$ from $\overline{H}$ and show $\overline{H}_{\mathbf{R}} \cdot g_0 = \pi^{-1}(M)$ ($g_0 \in \pi^{-1}(M)$), where π is the natural projection of G onto G/K . Furthermore, we construct a subgroup $\overline{\overline{H}}_{\mathbf{R}}$ of G from $\overline{H}_{\mathbf{R}}$ and show $\overline{\overline{H}}_{\mathbf{R}}(g_0K) = M$ ($g_0K \in M$). Thus the homogeneity of M follows.

Future plan of research By using (ii) of Corollary B and the equivalenceness of the above statements (I) and (II), we plan to investigate whether, for an irreducible C^ω -submanifold M in G/K of codimension greater than one, the following statement is true:

M is a principal orbit of a Hermann type action on G/K if and only if M is a curvature-adapted isoparametric submanifold with flat section admitting no non-Euclidean type focal point on $(G/K)(\infty)$.

Furthermore, we plan to investigate whether both the conditions of the curvature-adaptedness and the non-existenceness of non-Euclidean type focal point on $(G/K)(\infty)$ are indispensable in this statement. Note that this statement gives a submanifold geometrical characterization of a principal orbit of a Hermann type action.

2 Basic notions and facts

In this section, we recall basic notions introduced in [Koi1~3]. We first recall the notion of a complex equifocal submanifold introduced in [Koi1]. Let M be an immersed submanifold with abelian normal bundle (i.e., the sectional curvature for each 2-plane in the normal space is equal to zero) of in a symmetric space $N = G/K$ of non-compact type. Denote by A the shape tensor of M . Let $v \in T_x^\perp M$ and $X \in T_x M$ ($x = gK$). Denote by γ_v the geodesic in N with $\dot{\gamma}_v(0) = v$. The strongly M -Jacobi field Y along γ_v with $Y(0) = X$ (hence $Y'(0) = -A_v X$) is given by

$$Y(s) = (P_{\gamma_v|_{[0,s]}} \circ (D_{sv}^{co} - sD_{sv}^{si} \circ A_v))(X),$$

where $Y'(0) = \tilde{\nabla}_v Y$, $P_{\gamma_v|_{[0,s]}}$ is the parallel translation along $\gamma_v|_{[0,s]}$ and D_{sv}^{co} (resp. D_{sv}^{si}) is given by

$$\begin{aligned} D_{sv}^{co} &= g_* \circ \cos(\sqrt{-1}\text{ad}(sg_*^{-1}v)) \circ g_*^{-1} \\ \left(\text{resp. } D_{sv}^{si} &= g_* \circ \frac{\sin(\sqrt{-1}\text{ad}(sg_*^{-1}v))}{\sqrt{-1}\text{ad}(sg_*^{-1}v)} \circ g_*^{-1} \right). \end{aligned}$$

Here ad is the adjoint representation of the Lie algebra $\mathfrak{g}$ of G . All focal radii of M along γ_v are obtained as real numbers s_0 with $\text{Ker}(D_{s_0v}^{co} - s_0D_{s_0v}^{si} \circ A_v) \neq \{0\}$. So, we call a complex number z_0 with $\text{Ker}(D_{z_0v}^{co} - z_0D_{z_0v}^{si} \circ A_v^{\mathbb{C}}) \neq \{0\}$ a *complex focal radius of M along γ_v* and call $\dim \text{Ker}(D_{z_0v}^{co} - z_0D_{z_0v}^{si} \circ A_v^{\mathbb{C}})$ the *multiplicity* of the complex focal radius z_0 , where $D_{z_0v}^{co}$ (resp. $D_{z_0v}^{si}$) is a $\mathbb{C}$ -linear transformation of $(T_x N)^{\mathbb{C}}$ defined by

$$\begin{aligned} D_{z_0v}^{co} &= g_*^{\mathbb{C}} \circ \cos(\sqrt{-1}\text{ad}^{\mathbb{C}}(z_0g_*^{-1}v)) \circ (g_*^{\mathbb{C}})^{-1} \\ \left(\text{resp. } D_{z_0v}^{si} &= g_*^{\mathbb{C}} \circ \frac{\sin(\sqrt{-1}\text{ad}^{\mathbb{C}}(z_0g_*^{-1}v))}{\sqrt{-1}\text{ad}^{\mathbb{C}}(z_0g_*^{-1}v)} \circ (g_*^{\mathbb{C}})^{-1} \right), \end{aligned}$$

where $g_*^{\mathbb{C}}$ (resp. $\text{ad}^{\mathbb{C}}$) is the complexification of g_* (resp. ad). Here we note that, in the case where M is of class C^ω , complex focal radii along γ_v indicate the positions of focal points of the extrinsic complexification $M^{\mathbb{C}}(\hookrightarrow G^{\mathbb{C}}/K^{\mathbb{C}})$ of M along the complexified geodesic $\gamma_{\iota_*v}^{\mathbb{C}}$, where $G^{\mathbb{C}}/K^{\mathbb{C}}$ is the anti-Kaehlerian symmetric space associated with G/K and ι is the natural immersion of G/K into $G^{\mathbb{C}}/K^{\mathbb{C}}$. See the final paragraph of this

section about the definitions of G^c/K^c , $M^c(\hookrightarrow G^c/K^c)$ and $\gamma_{\iota_*v}^c$. Also, for a complex focal radius z_0 of M along γ_v , we call $z_0v \in (T_x^\perp M)^c$ a *complex focal normal vector of M at x* . Furthermore, assume that M has globally flat normal bundle (i.e., the normal holonomy group of M is trivial). Let $\tilde{v}$ be a parallel unit normal vector field of M . Assume that the number (which may be ∞) of distinct complex focal radii along $\gamma_{\tilde{v}_x}$ is independent of the choice of $x \in M$. Let $\{r_{i,x} | i = 1, 2, \dots\}$ be the set of all complex focal radii along $\gamma_{\tilde{v}_x}$, where $|r_{i,x}| < |r_{i+1,x}|$ or " $|r_{i,x}| = |r_{i+1,x}|$ & $\operatorname{Re} r_{i,x} > \operatorname{Re} r_{i+1,x}$ " or " $|r_{i,x}| = |r_{i+1,x}|$ & $\operatorname{Re} r_{i,x} = \operatorname{Re} r_{i+1,x}$ & $\operatorname{Im} r_{i,x} = -\operatorname{Im} r_{i+1,x} < 0$ ". Let r_i ($i = 1, 2, \dots$) be complex valued functions on M defined by assigning $r_{i,x}$ to each $x \in M$. We call these functions r_i ($i = 1, 2, \dots$) *complex focal radius functions for $\tilde{v}$* . We call $r_i\tilde{v}$ a *complex focal normal vector field for $\tilde{v}$* . If, for each parallel unit normal vector field $\tilde{v}$ of M , the number of distinct complex focal radii along $\gamma_{\tilde{v}_x}$ is independent of the choice of $x \in M$, each complex focal radius function for $\tilde{v}$ is constant on M and it has constant multiplicity, then we call M a *complex equifocal submanifold*. Let $\phi : H^0([0, 1], \mathfrak{g}) \rightarrow G$ be the parallel transport map for G , that is, $\phi(u) := g_u(1)$ ($u \in H^0([0, 1], \mathfrak{g})$), where g_u is the element of $H^1([0, 1], G)$ with $g_u(0) = e$ and $g_{u*}^{-1}g'_u = u$. See Section 4 of [Koi1] about the definitions of $H^0([0, 1], \mathfrak{g})$ and $H^1([0, 1], G)$. Let $\pi : G \rightarrow G/K$ be the natural projection. It follows from Theorem 1 of [Koi2] that, M is complex equifocal if and only if each component of $(\pi \circ \phi)^{-1}(M)$ is complex isoparametric. See Section 2 of [Koi1] about the definition of a complex isoparametric submanifold. In particular, if each component of $(\pi \circ \phi)^{-1}(M)$ is proper complex isoparametric (i.e., complex isoparametric and, for each unit normal vector v , the complexified shape operator A_v^c is diagonalizable with respect to a pseudo-orthonormal base), then we call M a *proper complex equifocal submanifold*.

Next we recall the notion of an infinite dimensional proper anti-Kaehlerian isoparametric submanifold. Let M be an anti-Kaehlerian Fredholm submanifold in an infinite dimensional anti-Kaehlerian space V and A be the shape tensor of M . See [Koi2] about the definitions of an infinite dimensional anti-Kaehlerian space and anti-Kaehlerian Fredholm submanifold in the space. Denote by the same symbol J the complex structures of M and V . Fix a unit normal vector v of M . If there exists $X(\neq 0) \in TM$ with $A_v X = aX + bJX$, then we call the complex number $a + b\sqrt{-1}$ a *J -eigenvalue of A_v* (or a *complex principal curvature of direction v*) and call X a *J -eigenvector for $a + b\sqrt{-1}$* . Also, we call the space of all J -eigenvectors for $a + b\sqrt{-1}$ a *J -eigenspace for $a + b\sqrt{-1}$* . The J -eigenspaces are orthogonal to one another and each J -eigenspace is J -invariant. We call the set of all J -eigenvalues of A_v the *J -spectrum of A_v* and denote it by $\operatorname{Spec}_J A_v$. The set $\operatorname{Spec}_J A_v \setminus \{0\}$ is described as follows:

$$\operatorname{Spec}_J A_v \setminus \{0\} = \{\lambda_i | i = 1, 2, \dots\}$$

$$\left(\begin{array}{l} |\lambda_i| > |\lambda_{i+1}| \text{ or } "|\lambda_i| = |\lambda_{i+1}| \text{ \& } \operatorname{Re} \lambda_i > \operatorname{Re} \lambda_{i+1}" \\ \text{or } "|\lambda_i| = |\lambda_{i+1}| \text{ \& } \operatorname{Re} \lambda_i = \operatorname{Re} \lambda_{i+1} \text{ \& } \operatorname{Im} \lambda_i = -\operatorname{Im} \lambda_{i+1} > 0" \end{array} \right).$$

Also, the J -eigenspace for each J -eigenvalue of A_v other than 0 is of finite dimension. We call the J -eigenvalue λ_i the i -th complex principal curvature of direction v . Assume that M has globally flat normal bundle. Fix a parallel normal vector field $\tilde{v}$ of M . Assume that the number (which may be ∞) of distinct complex principal curvatures of direction $\tilde{v}_x$ is independent of the choice of $x \in M$. Then we can define functions $\tilde{\lambda}_i$ ($i = 1, 2, \dots$) on M by assigning the i -th complex principal curvature of direction $\tilde{v}_x$ to each $x \in M$. We call this function $\tilde{\lambda}_i$ the i -th complex principal curvature function of direction $\tilde{v}$. We consider the following condition:

(AKI) For each parallel normal vector field $\tilde{v}$, the number of distinct complex principal curvatures of direction $\tilde{v}_x$ is independent of the choice of $x \in M$, each complex principal curvature function of direction $\tilde{v}$ is constant on M and it has constant multiplicity.

If M satisfies this condition (AKI), then we call M an *anti-Kaehlerian isoparametric submanifold*. Let $\{e_i\}_{i=1}^\infty$ be an orthonormal system of $T_x M$. If $\{e_i\}_{i=1}^\infty \cup \{Je_i\}_{i=1}^\infty$ is an orthonormal base of $T_x M$, then we call $\{e_i\}_{i=1}^\infty$ a *J -orthonormal base*. If there exists a J -orthonormal base consisting of J -eigenvectors of A_v , then A_v is said to be *diagonalized with respect to the J -orthonormal base*. If M is anti-Kaehlerian isoparametric and, for each $v \in T^\perp M$, the shape operator A_v is diagonalized with respect to a J -orthonormal base, then we call M a *proper anti-Kaehlerian isoparametric submanifold*. For arbitrary two unit normal vector v_1 and v_2 of a proper anti-Kaehlerian isoparametric submanifold, the shape operators A_{v_1} and A_{v_2} are simultaneously diagonalized with respect to a J -orthonormal base. Assume that M is a proper anti-Kaehlerian isoparametric submanifold. Let $\{E_i \mid i \in I\}$ be the family of distributions on M such that, for each $x \in M$, $\{(E_i)_x \mid i \in I\}$ is the set of all common J -eigenspaces of A_v 's ($v \in T_x^\perp M$). The relation $TM = \overline{\bigoplus_{i \in I} E_i}$ holds.

Let λ_i ($i \in I$) be the section of $(T^\perp M)^* \otimes \mathbf{C}$ such that $A_v = \operatorname{Re} \lambda_i(v) \operatorname{id} + \operatorname{Im} \lambda_i(v) J$ on $(E_i)_{\pi(v)}$ for each $v \in T^\perp M$, where π is the bundle projection of $T^\perp M$. We call λ_i ($i \in I$) *complex principal curvatures* of M and call distributions E_i ($i \in I$) *complex curvature distributions* of M . It is shown that there uniquely exists a normal vector field n_i of M with $\lambda_i(\cdot) = \langle n_i, \cdot \rangle - \sqrt{-1} \langle Jn_i, \cdot \rangle$ (see Lemma 5 of [Koi2]). We call n_i ($i \in I$) the *complex curvature normals* of M . Note that n_i is parallel with respect to the normal connection $\nabla^\perp$. Similarly we can define a (finite dimensional) proper anti-Kaehlerian isoparametric submanifold in a finite dimensional anti-Kaehlerian space, its complex principal curvatures, its complex curvature distributions and its complex curvature normals. Set $l_i^x := (\lambda_i)_x^{-1}(1)$. In [Koi2], it has been shown that the focal set of (M, x) is equal to $\bigcup_{i \in I} l_i^x$. Denote by T_i^x the complex reflection of order 2 with respect to the complex hyperplane l_i^x of $T_x^\perp M$ (i.e., the rotation of angle π having l_i^x as the axis), which is an affine transformation of $T_x^\perp M$. Let W_x be the group generated by T_i^x 's ($i \in I$). According to Proposition 3.7 of [Koi4], W_x is discrete and it is independent of the choice of $x \in M$ (up to group isomorphism). Hence we simply denote it by W . We call this group W the *complex Coxeter group associated*

with M . According to Lemma 3.8 of [Koi4], W is decomposable (i.e., it is decomposed into a non-trivial product of two discrete complex reflection groups) if and only if there exist two J -invariant linear subspaces $P_1 (\neq \{0\})$ and $P_2 (\neq \{0\})$ of $T_x^\perp M$ such that $T_x^\perp M = P_1 \oplus P_2$ (orthogonal direct sum), $P_1 \cup P_2$ contains all complex curvature normals of M at x and that P_i ($i = 1, 2$) contains at least one complex curvature normal of M at x .

Next we recall the notion of the extrinsic complexification of a complete C^ω -submanifold in a symmetric space of non-compact type which was introduced in [Koi2]. First we recall the notions of an anti-Kaehlerian symmetric space associated with a symmetric space of non-compact type and an aks-representation. Let J be a parallel complex structure on an even dimensional pseudo-Riemannian manifold $(M, \langle \cdot, \cdot \rangle)$ of half index. If $\langle JX, JY \rangle = -\langle X, Y \rangle$ holds for every $X, Y \in TM$, then $(M, \langle \cdot, \cdot \rangle, J)$ is called an *anti-Kaehlerian manifold*. Let G/K be a symmetric space of non-compact type and $(\mathfrak{g}, \sigma)$ be its orthogonal symmetric Lie algebra. Let $\mathfrak{g} = \mathfrak{f} + \mathfrak{p}$ be the Cartan decomposition associated with a symmetric pair (G, K) . Note that $\mathfrak{f}$ is the Lie algebra of K and $\mathfrak{p}$ is identified with the tangent space $T_e K(G/K)$, where e is the identity element of G . Let $\langle \cdot, \cdot \rangle$ be the $\text{Ad}_G(G)$ -invariant non-degenerate inner product of $\mathfrak{g}$ inducing the Riemannian metric of G/K , where Ad_G is the adjoint representation of G . Let $G^\mathbb{C}$ (resp. $K^\mathbb{C}$) be the complexification of G (resp. K). Without loss of generality, we may assume that $K^\mathbb{C}$ is connected and that $G^\mathbb{C}$ is simply connected. The 2-multiple of the real part $\text{Re}\langle \cdot, \cdot \rangle^\mathbb{C}$ of $\langle \cdot, \cdot \rangle^\mathbb{C}$ is the Killing form of $\mathfrak{g}^\mathbb{C}$ regarded as a real Lie algebra. The restriction $2\text{Re}\langle \cdot, \cdot \rangle^\mathbb{C}|_{\mathfrak{p}^\mathbb{C} \times \mathfrak{p}^\mathbb{C}}$ is an $\text{Ad}(K^\mathbb{C})$ -invariant non-degenerate inner product of $\mathfrak{p}^\mathbb{C}$ ($= T_{eK^\mathbb{C}}(G^\mathbb{C}/K^\mathbb{C})$). Denote by $\langle \cdot, \cdot \rangle^A$ the $G^\mathbb{C}$ -invariant pseudo-Riemannian metric on $G^\mathbb{C}/K^\mathbb{C}$ induced from $2\text{Re}\langle \cdot, \cdot \rangle^\mathbb{C}|_{\mathfrak{p}^\mathbb{C} \times \mathfrak{p}^\mathbb{C}}$. Define an almost complex structure J_0 of $\mathfrak{p}^\mathbb{C}$ by $J_0 X = \sqrt{-1}X$ ($X \in \mathfrak{p}^\mathbb{C}$). It is clear that J_0 is $\text{Ad}(K^\mathbb{C})$ -invariant. Denote by $\tilde{J}$ the $G^\mathbb{C}$ -invariant almost complex structure on $G^\mathbb{C}/K^\mathbb{C}$ induced from J_0 . It is shown that $(G^\mathbb{C}/K^\mathbb{C}, \langle \cdot, \cdot \rangle^A, \tilde{J})$ is an anti-Kaehlerian manifold and a (semi-simple) pseudo-Riemannian symmetric space. We call this anti-Kaehlerian manifold an *anti-Kaehlerian symmetric space associated with G/K* and simply denote it by $G^\mathbb{C}/K^\mathbb{C}$. The action $\text{Ad}_{G^\mathbb{C}}(K^\mathbb{C})$ on $\mathfrak{g}^\mathbb{C}$ preserves $\mathfrak{p}^\mathbb{C}$ invariantly, where $\text{Ad}_{G^\mathbb{C}}$ is the adjoint representation of $G^\mathbb{C}$. Define a representation ρ of $K^\mathbb{C}$ on $\mathfrak{p}^\mathbb{C}$ by $\rho(k)(X) := \text{Ad}_{G^\mathbb{C}}(k)(X)$ ($k \in K^\mathbb{C}, X \in \mathfrak{p}^\mathbb{C}$). If $G^\mathbb{C}/K^\mathbb{C}$ is irreducible, then we call this representation ρ an *aks-representation (associated with $G^\mathbb{C}/K^\mathbb{C}$)*. Let X_0 be a semi-simple element of $\mathfrak{p}^\mathbb{C}$, where the semi-simplicity means that the complexification of $\text{ad}_{\mathfrak{g}^\mathbb{C}}(X_0)$ is diagonalizable. If the orbit $\rho(K^\mathbb{C}) \cdot X_0$ is principal, then it is a (finite dimensional) proper anti-Kaehlerian isoparametric submanifold (see Lemma 3.5.3). Let N be a complete C^ω -Riemannian manifold. The notion of the adapted complex structure on a neighborhood U of the 0-section of the tangent bundle TN is defined as the complex structure (on U) such that, for each geodesic $\gamma : \mathbf{R} \rightarrow N$, the restriction of its differential $\gamma_* : T\mathbf{R} = \mathbf{C} \rightarrow TN$ to $\gamma_*^{-1}(U)$ is holomorphic. We take U as largely as possible under the condition that $U \cap T_x N$ is a star-shaped

neighborhood of 0_x for each $x \in N$, where 0_x is the zero vector of $T_x N$. If N is of non-negative curvature, then we have $U = TN$. Also, if all sectional curvatures of N are bigger than or equal to c ($c < 0$), then U contains the ball bundle $T^r N := \{X \in TN \mid \|X\| < r\}$ of radius $r := \frac{\pi}{2\sqrt{-c}}$. In detail, see [S1,2]. Denote by J_A the adapted complex structure on U . The complex manifold (U, J_A) is interpreted as the complexification of N . We denote (U, J_A) by N^c and call it the complexification of N , where we note that N^c is given no pseudo-Riemannian metric. In particular, in case of $N = \mathbf{R}^m$ (the Euclidean space), we have $(U, J_A) = \mathbf{C}^m$. Also, in the case where N is a symmetric space G/K of non-compact type, there exists the holomorphic diffeomorphism δ of (U, J_A) onto an open subset of G^c/K^c . Let M be an immersed (complete) C^ω -submanifold in G/K . Denote by f its immersion. Let M^c be the complexification of M (defined as above). We shall state the definition of the complexification $f^c : M^c \rightarrow G^c/K^c$ of f , where we shrink M^c to a neighborhood of the 0-section of TM if necessary. For its purpose, we first state the definition of the complexification of a C^ω -curve $\alpha : \mathbf{R} \rightarrow G/K$. Let $\mathfrak{g} = \mathfrak{f} + \mathfrak{p}$ be the Cartan decomposition associated with G/K and $W : \mathbf{R} \rightarrow \mathfrak{p}$ be the curve in $\mathfrak{p}$ with $(\exp W(t))K = \alpha(t)$ ($t \in \mathbf{R}$), where we note that W is uniquely determined because G/K is of non-compact type. Since α is of class C^ω , so is also W . Let $W^c : D \rightarrow \mathfrak{p}^c$ ($D : \text{a neighborhood of } \mathbf{R} \text{ in } \mathbf{C}$) be the holomorphic extension of W . We define the complexification $\alpha^c : D \rightarrow G^c/K^c$ of α by $\alpha^c(z) = (\exp W^c(z))K^c$. It is shown that this complexification of a C^ω -curve in G/K is a holomorphic curve in G^c/K^c . By using this complexification of a C^ω -curve in G/K , we define the complexification $f^c : M^c \rightarrow G^c/K^c$ of f by $f^c(X) := (f \circ \gamma_X^M)^c(\sqrt{-1})$ ($X \in M^c \subset TM$), where γ_X^M is the geodesic in M with $\dot{\gamma}_X^M(0) = X$. Here we shrink M^c to a neighborhood of the 0-section of TM if necessary in order to assure that $\sqrt{-1}$ belongs to the domain of $(f \circ \gamma_X^M)^c$ for each $X \in M^c$. It is shown that the map $f^c : M^c \rightarrow G^c/K^c$ is holomorphic and that the restriction of f^c to a neighborhood U' of the 0-section of TM is an immersion, where we take U' as largely as possible. Denote by M^c this neighborhood U' newly. Give M^c the pseudo-Riemannian metric induced from that of G^c/K^c by f^c . Then M^c is an anti-Kaehlerian submanifold in G^c/K^c immersed by f^c . We call this anti-Kaehlerian submanifold M^c immersed by f^c the *extrinsic complexification* of the submanifold M . In [Koi2] and [Koi6], we constructed the complete extension of the extrinsic complexification in different methods in the case where M is proper complex equifocal. We also denote this complete extension by the same symbol M^c . Here we note that the extrinsic complexification of a C^ω -pseudo-Riemannian submanifold in a general pseudo-Riemannian manifold has recently defined in [Koi7].

Let $\langle \cdot, \cdot \rangle$ be the $\text{Ad}(G)$ -invariant non-degenerate symmetric bilinear form of $\mathfrak{g}$ inducing the metric of G/K . The Cartan decomposition $\mathfrak{g} = \mathfrak{f} \oplus \mathfrak{p}$ is an orthogonal time-space decomposition of $\mathfrak{g}$ with respect to $\langle \cdot, \cdot \rangle$ in the sense of [Koi1]. Set $\langle \cdot, \cdot \rangle^A := 2\text{Re}\langle \cdot, \cdot \rangle^c$, where $\langle \cdot, \cdot \rangle^c$ is the complexification of $\langle \cdot, \cdot \rangle$ (which is a $\mathbf{C}$ -bilinear form of $\mathfrak{g}^c$). The $\mathbf{R}$ -bilinear form $\langle \cdot, \cdot \rangle^A$ on $\mathfrak{g}^c$ regarded as a real Lie algebra induces a bi-invariant pseudo-

Riemannian metric of G^c and a G^c -invariant anti-Kaehlerian metric on G^c/K^c . It is clear that $\mathfrak{g}^c = (\mathfrak{f} + \sqrt{-1}\mathfrak{p}) \oplus (\sqrt{-1}\mathfrak{f} + \mathfrak{p})$ is an orthogonal time-space decomposition of $\mathfrak{g}^c$ with respect to $\langle \cdot, \cdot \rangle^A$. For simplicity, set $\mathfrak{g}_-^c := \mathfrak{f} + \sqrt{-1}\mathfrak{p}$, $\mathfrak{g}_+^c := \sqrt{-1}\mathfrak{f} + \mathfrak{p}$. Set $\langle \cdot, \cdot \rangle_{\mathfrak{g}_\pm^c}^A := -\pi_{\mathfrak{g}_-^c}^* \langle \cdot, \cdot \rangle^A + \pi_{\mathfrak{g}_\pm^c}^* \langle \cdot, \cdot \rangle^A$, where $\pi_{\mathfrak{g}_-^c}$ (resp. $\pi_{\mathfrak{g}_+^c}$) is the projection of $\mathfrak{g}^c$ onto $\mathfrak{g}_-^c$ (resp. $\mathfrak{g}_+^c$). Let $H^0([0, 1], \mathfrak{g}^c)$ be the space of all L^2 -integrable paths $u : [0, 1] \rightarrow \mathfrak{g}^c$ with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{g}_\pm^c}^A$ and $H^0([0, 1], \mathfrak{g}_-^c)$ (resp. $H^0([0, 1], \mathfrak{g}_+^c)$) be the space of all L^2 -integrable paths $u : [0, 1] \rightarrow \mathfrak{g}_-^c$ (resp. $u : [0, 1] \rightarrow \mathfrak{g}_+^c$) with respect to $-\langle \cdot, \cdot \rangle_{\mathfrak{g}_-^c \times \mathfrak{g}_-^c}^A$ (resp. $\langle \cdot, \cdot \rangle_{\mathfrak{g}_+^c \times \mathfrak{g}_+^c}^A$). It is clear that $H^0([0, 1], \mathfrak{g}^c) = H^0([0, 1], \mathfrak{g}_-^c) \oplus H^0([0, 1], \mathfrak{g}_+^c)$. Define a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle_0^A$ of $H^0([0, 1], \mathfrak{g}^c)$ by $\langle u, v \rangle_0^A := \int_0^1 \langle u(t), v(t) \rangle^A dt$. It is easy to show that the decomposition $H^0([0, 1], \mathfrak{g}^c) = H^0([0, 1], \mathfrak{g}_-^c) \oplus H^0([0, 1], \mathfrak{g}_+^c)$ is an orthogonal time-space decomposition with respect to $\langle \cdot, \cdot \rangle_0^A$. For simplicity, set $H_\pm^{0,c} := H^0([0, 1], \mathfrak{g}_\pm^c)$ and $\langle \cdot, \cdot \rangle_{0, H_\pm^{0,c}}^A := -\pi_{H_-^{0,c}}^* \langle \cdot, \cdot \rangle_0^A + \pi_{H_\pm^{0,c}}^* \langle \cdot, \cdot \rangle_0^A$, where $\pi_{H_-^{0,c}}$ (resp. $\pi_{H_+^{0,c}}$) is the projection of $H^0([0, 1], \mathfrak{g}^c)$ onto $H_-^{0,c}$ (resp. $H_+^{0,c}$). It is clear that $\langle u, v \rangle_{0, H_\pm^{0,c}}^A = \int_0^1 \langle u(t), v(t) \rangle_{\mathfrak{g}_\pm^c}^A dt$ ($u, v \in H^0([0, 1], \mathfrak{g}^c)$). Hence $(H^0([0, 1], \mathfrak{g}^c), \langle \cdot, \cdot \rangle_{0, H_\pm^{0,c}}^A)$ is a Hilbert space, that is, $(H^0([0, 1], \mathfrak{g}^c), \langle \cdot, \cdot \rangle_0^A)$ is a pseudo-Hilbert space in the sense of [Koi1]. Let J be the endomorphism of $\mathfrak{g}^c$ defined by $JX = \sqrt{-1}X$ ($X \in \mathfrak{g}^c$). Denote by the same symbol J the bi-invariant almost complex structure of G^c induced from J . Define the endomorphism $\tilde{J}$ of $H^0([0, 1], \mathfrak{g}^c)$ by $\tilde{J}u = \sqrt{-1}u$ ($u \in H^0([0, 1], \mathfrak{g}^c)$). Since $\tilde{J}H_\pm^{0,c} = H_\mp^{0,c}$ and $\langle \tilde{J}u, \tilde{J}v \rangle_0^A = -\langle u, v \rangle_0^A$ ($u, v \in H^0([0, 1], \mathfrak{g}^c)$), the space $(H^0([0, 1], \mathfrak{g}^c), \langle \cdot, \cdot \rangle_0^A, \tilde{J})$ is an anti-Kaehlerian space. Let $H^1([0, 1], \mathfrak{g}^c)$ be a Hilbert subspace of $H^0([0, 1], \mathfrak{g}^c)$ consisting of all absolutely continuous paths $u : [0, 1] \rightarrow \mathfrak{g}^c$ such that the weak derivative u' of u is squared integrable (with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{g}_\pm^c}^A$). Also, let $H^1([0, 1], G^c)$ be the Hilbert Lie group of all absolutely continuous paths $g : [0, 1] \rightarrow G^c$ such that the weak derivative g' of g is squared integrable (with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{g}_\pm^c}^A$), that is, $g_*^{-1}g' \in H^0([0, 1], \mathfrak{g}^c)$. Let $\phi^c : H^0([0, 1], \mathfrak{g}^c) \rightarrow G^c$ be the parallel transport map for G^c , that is, $\phi^c(u) := g_u(1)$ ($u \in H^0([0, 1], \mathfrak{g}^c)$), where g_u is the element of $H^1([0, 1], G^c)$ with $g_u(0) = e$ and $g_{u*}^{-1}g'_u = u$. This map ϕ^c is an anti-Kaehlerian submersion. Set $P(G^c, e \times G^c) := \{g \in H^1([0, 1], G^c) \mid g(0) = e\}$ and $\Omega_e(G^c) := \{g \in H^1([0, 1], G^c) \mid g(0) = g(1) = e\}$. The group $H^1([0, 1], G^c)$ acts on $H^0([0, 1], \mathfrak{g}^c)$ by gauge transformations, that is,

$$g * u := \text{Ad}(g)u - g'g_*^{-1} \quad (g \in H^1([0, 1], G^c), u \in H^0([0, 1], \mathfrak{g}^c)).$$

It is shown that the following facts hold:

- (i) The above action of $H^1([0, 1], G^c)$ on $H^0([0, 1], \mathfrak{g}^c)$ is isometric,
- (ii) The above action of $P(G^c, e \times G^c)$ on $H^0([0, 1], \mathfrak{g}^c)$ is transitive and free,
- (iii) $\phi^c(g * u) = (L_{g(0)} \circ R_{g(1)}^{-1})(\phi^c(u))$ for $g \in H^1([0, 1], G^c)$ and $u \in H^0([0, 1], \mathfrak{g}^c)$,
- (iv) $\phi^c : H^0([0, 1], \mathfrak{g}^c) \rightarrow G^c$ is regarded as a $\Omega_e(G^c)$ -bundle.

(v) If $\phi^c(u) = (L_{x_0} \circ R_{x_1}^{-1})(\phi^c(v))$ ($u, v \in H^0([0, 1], \mathfrak{g}^c)$, $x_0, x_1 \in G^c$), then there exists $g \in H^1([0, 1], G^c)$ such that $g(0) = x_0$, $g(1) = x_1$ and $u = g * v$. In particular, it follows that each $u \in H^0([0, 1], \mathfrak{g}^c)$ is described as $u = g * \hat{0}$ in terms of some $g \in P(G^c, G^c \times e)$. Let $\pi^c : G^c \rightarrow G^c/K^c$ be the natural projection. It is shown that M is proper complex equifocal if and only if $\widetilde{M}^c := (\pi^c \circ \phi^c)^{-1}(M^c)$ is proper anti-Kaehlerian isoparametric. The focal set of M^c at $x(\in M)$ is equal to the image by the normal exponential map (of M^c) of the focal set (which consists of complex hyperplanes in $T_u^\perp \widetilde{M}^c$) of $\widetilde{M}^c$ at $u \in \widetilde{M}^c \cap (\pi \circ \phi)^{-1}(x)$ under the identification of $T_u^\perp \widetilde{M}^c$ with $T_x^\perp M^c$. From this reason, we call the above complex Coxeter group associated with $\widetilde{M}^c$ the *complex Coxeter group associated with M* .

3 Homogeneity of the lifted submanifold

Let M be an irreducible proper complex equifocal submanifold of codimension greater than one in a symmetric space G/K of non-compact type and $\widetilde{M}^c$ be the lifted submanifold of the complexification of M to $H^0([0, 1], \mathfrak{g}^c)$. In this section, we shall prove the homogeneity of $\widetilde{M}^c$. First we shall recall the generalized Chow's theorem, which was proved in [HL2]. Let N be a (connected) Hilbert manifold and $\mathcal{D}$ be a set of local (smooth) vector fields which are open sets of N . If two points x and y of N can be connected by a piecewise smooth curve each of whose smooth segments is an integral curve of a local smooth vector field belonging to $\mathcal{D}$, then we say that x and y are $\mathcal{D}$ -equivalent and we denote this fact by $x \sim_{\mathcal{D}} y$. Let $\Omega_{\mathcal{D}}(x) := \{y \in N \mid y \sim_{\mathcal{D}} x\}$. The set $\Omega_{\mathcal{D}}(x)$ is called the *set of reachable points of $\mathcal{D}$ starting from x* . Let $\mathcal{D}^*$ be the set of local smooth vector fields on open sets of N which is generated by $\mathcal{D}$ in the following sense: $\mathcal{D} \subset \mathcal{D}^*$, $\mathcal{D}^*$ contains the zero vector field and, for any $X, Y \in \mathcal{D}^*$ and any $a, b \in \mathbf{R}$, $aX + bY$ and $[X, Y]$ (which are defined on the intersection of the domains of X and Y) also belong to $\mathcal{D}^*$. For each $x \in N$, let $\mathcal{D}^*(x) := \{X_x \mid X \in \mathcal{D}^* \text{ s.t. } x \in \text{Dom}(X)\}$. Then the following generalized Chow's theorem holds.

Theorem 3.1([HL2]) *If $\overline{\mathcal{D}^*(x)} = T_x N$ for each $x \in N$, $\overline{\Omega_{\mathcal{D}}(x)} = N$ holds for each $x \in N$, where $\overline{(\cdot)}$ implies the closure of $(\cdot)$.*

For simplicity, we set $V := H^0([0, 1], \mathfrak{g}^c)$. Denote by A the shape tensor of $\widetilde{M}^c$. Let $\{E_i \mid i \in I\} \cup \{E_0\}$ be the set of all complex curvature distributions of $\widetilde{M}^c$, where E_0 is one defined by $(E_0)_u := \bigcap_{v \in T_u^\perp \widetilde{M}^c} \text{Ker } \widetilde{A}_v$ ($u \in \widetilde{M}^c$). Also, let λ_i and n_i be the complex principal curvature and the complex curvature normal corresponding to E_i , respectively. Fix $u_0 \in \widetilde{M}^c$. Denote by l_i the complex hyperplane $(\lambda_i)_{u_0}^{-1}(1)$ of $T_{u_0}^\perp \widetilde{M}^c$. Let $Q(u_0)$ be the set of all points of $\widetilde{M}^c$ connected with u_0 by a piecewise smooth curve in $\widetilde{M}^c$ each of whose

smooth segments is contained in some complex curvature sphere (which may depend on the smooth segment). By using the above generalized Chow's theorem, we shall show the following fact.

Proposition 3.2. *The set $Q(u_0)$ is dense in $\widetilde{M}^c$.*

Proof. Let $\mathcal{D}_E$ be the set of all local (smooth) tangent vector fields on open sets of $\widetilde{M}^c$ which is tangent to some E_i ($i \neq 0$) at each point of the domain. Define $\Omega_{\mathcal{D}_E}(u_0)$, $\mathcal{D}_E^*$ and $\mathcal{D}_E^*(u_0)$ as above. By imitating the proof of Proposition 5.8 of [HL1], it is shown that $\overline{\mathcal{D}_E^*(u)} = T_u \widetilde{M}^c$ for each $u \in \widetilde{M}^c$. Hence, $\overline{\Omega_{\mathcal{D}_E}(u_0)} = \widetilde{M}^c$ follows from Theorem 3.1. It is clear that $\Omega_{\mathcal{D}_E}(u_0) = Q(u_0)$. Therefore we obtain $\overline{Q(u_0)} = \widetilde{M}^c$. q.e.d.

For each complex affine subspace P of $T_{u_0}^\perp \widetilde{M}^c$, define I_P by

$$I_P := \begin{cases} \{i \in I \mid (n_i)_{u_0} \in P\} & (0 \notin P) \\ \{i \in I \mid (n_i)_{u_0} \in P\} \cup \{0\} & (0 \in P). \end{cases}$$

It is easy to show that I_P is finite. Define a distribution D_P on $\widetilde{M}^c$ by $D_P := \bigoplus_{i \in I_P} E_i$. It is shown that $(\bigcap_{i \in I_P \setminus \{0\}} l_i) \setminus (\bigcup_{i \in I \setminus I_P} l_i) \neq \emptyset$. Take $v_0 \in (\bigcap_{i \in I_P \setminus \{0\}} l_i) \setminus (\bigcup_{i \in I \setminus I_P} l_i)$. Let v be a parallel normal vector field on $\widetilde{M}^c$ with $v_{u_0} = v_0$. Hence we note that v is a complex focal normal vector field of $\widetilde{M}^c$. Let f_v be the focal map for v (i.e., the end point map for v), F_v be the focal submanifold for v (i.e., $F_v = f_v(\widetilde{M}^c)$) and $L_u^{D_P}$ be the leaf of D_P through $u \in \widetilde{M}^c$. If $0 \notin P$, then we have $L_u^{D_P} = f_v^{-1}(f_v(u))$. According to the homogeneous slice theorem for the complete complexification of a proper complex equifocal submanifold in [Koi6], we have the following homogeneous slice theorem for $\widetilde{M}^c$.



Theorem 3.3. *If $0 \notin P$, then the leaf $L_u^{D_P}$ is a principal orbit of the direct sum representation of aks-representations on $T_{f_v(u)}^\perp F_v$.*

Proof. Let $u_1 := f_v(u)$, $\overline{F}_v := (\pi^c \circ \phi^c)(F_v)$ and $\overline{v} := (\pi^c \circ \phi^c)_*(v)$, which is well-defined because v is projectable. It is shown that $\overline{v}$ is a focal normal vector field of M^c and that $\overline{F}_v$ is the focal submanifold of M^c corresponding to $\overline{v}$. Denote by $f_{\overline{v}}$ the focal map for $\overline{v}$ and set $\overline{u}_1 := (\pi^c \circ \phi^c)(u_1)$. Set $L := f_v^{-1}(u_1)$ and $\overline{L} := f_{\overline{v}}^{-1}(\overline{u}_1)$, which are leaves of the focal distributions corresponding to v and $\overline{v}$, respectively. According to Theorem A of [Koi6], $\overline{L}$ is the image of a principal orbit of the direct sum representation of aks-representations on $T_{\overline{u}_1}^\perp \overline{F}_v$ by the normal exponential map $\exp_{\overline{u}_1}^\perp$ of $\overline{F}_v$ at $\overline{u}_1$. On the other hand, under the identification of $T_{\overline{u}_1}^\perp \overline{F}_v$ with $T_{u_1}^\perp F_v$, $\overline{L}$ is the image of L by $\exp_{\overline{u}_1}^\perp$. Hence it follows that L is a principal orbit of an aks-representation on $T_{u_1}^\perp F_v$. Since $0 \notin P$ by the assumption, we have $L_u^{D_P} = L$. Therefore the statement of this theorem follows. q.e.d.

Set $(W_P)_u := u + (D_P)_u \oplus \text{Span}\{(n_i)_u \mid i \in I_P \setminus \{0\}\}$ ($u \in \widetilde{M}^c$). Let $\gamma : [0, 1] \rightarrow \widetilde{M}^c$ be a piecewise smooth curve. In the sequel, we assume that the domains of all piecewise smooth curves are equal to $[0, 1]$. If $\dot{\gamma}(t) \perp (D_P)_{\gamma(t)}$ for each $t \in [0, 1]$, then γ is said to be *horizontal with respect to D_P* (or *D_P -horizontal*). Let β_i ($i = 1, 2$) be curves in $\widetilde{M}^c$. If $L_{\beta_1(t)}^{D_P} = L_{\beta_2(t)}^{D_P}$ for each $t \in [0, 1]$, then β_1 and β_2 are said to be *parallel*. By imitating the proof of Proposition 1.1 in [HL2], we can show the following fact.

Lemma 3.4. *For each D_P -horizontal curve γ , there exists an one-parameter family $\{h_{\gamma,t}^{D_P} \mid 0 \leq t \leq 1\}$ of holomorphic isometries $h_{\gamma,t}^{D_P} : (W_P)_{\gamma(0)} \rightarrow (W_P)_{\gamma(t)}$ such that $h_{\gamma,t}^{D_P}(L_{\gamma(0)}^{D_P}) = L_{\gamma(t)}^{D_P}$ and that, for each $u \in L_{\gamma(0)}^{D_P}$, $t \mapsto h_{\gamma,t}^{D_P}(u)$ is a D_P -horizontal curve parallel to γ .*

Proof. First we consider the case of $0 \notin P$. Take $v_0 \in \bigcap_{i \in I_P} l_i \setminus (\bigcup_{i \in I \setminus I_P} l_i)$. Let v be the parallel normal vector field of $\widetilde{M}^c$ with $v_{u_0} = v_0$. Let $\bar{\gamma} := f_v \circ \gamma$. Define a map $h_t : (W_P)_{\gamma(0)} \rightarrow V$ by $h_t(u) := \bar{\gamma}(t) + \tau_{\bar{\gamma}}^\perp(\bar{\gamma}(0)u)$ ($u \in (W_P)_{\gamma(0)}$), where $\tau_{\bar{\gamma}}^\perp$ is the parallel translation along $\bar{\gamma}$ with respect to the normal connection of F_v . Then it is shown that $\{h_t \mid 0 \leq t \leq 1\}$ is the desired one-parameter family (see Fig. 1). Next we consider the case of $0 \in P$. Take $v_0 \in \bigcap_{i \in I_P} (\lambda_i)_{u_0}^{-1}(0) \setminus (\bigcup_{i \in I \setminus I_P} (\lambda_i)_{u_0}^{-1}(0))$. Let v be the parallel normal vector field of $\widetilde{M}^c$ with $v_{u_0} = v_0$. We define a map $\nu : \widetilde{M}^c \rightarrow S^\infty(1)$ by $\nu(u) := v_u$ ($u \in \widetilde{M}^c$), where $S^\infty(1)$ is the unit hypersphere of V . Then we have $\nu_{*u} = -A_{v_u}$ ($u \in \widetilde{M}^c$), where A is the shape tensor of $\widetilde{M}^c$. If $i \in I_P$, then we have $\nu_{*u}((E_i)_u) = -\langle (n_i)_u, v_u \rangle (E_i)_u = \{0\}$ and, if $i \notin I_P$, then we have $\nu_{*u}((E_i)_u) = -\langle (n_i)_u, v_u \rangle (E_i)_u = (E_i)_u$. Hence we have $\text{Ker } \nu_{*u} = (D_P)_u$. Therefore D_P is integrable and it gives a foliation on $\widetilde{M}^c$. Denote by $\mathfrak{F}_P$ this foliation and $D_P^\perp$ the orthogonal complementary distribution of $\mathfrak{F}_P$. Let U be a neighborhood of $\gamma(0)$ in $L_{\gamma(0)}^{D_P}$ where an element of holonomy along γ with respect to $(\mathfrak{F}_D, D_P^\perp)$ is defined. See [BH] about the definition of an element of holonomy. Let Δ be a fundamental domain containing u_0 of the complex Coxeter group of $\widetilde{M}^c$ at $u_0 (\in \widetilde{M}^c)$. Denote by Δ_u a domain of $T_u^\perp \widetilde{M}^c$ given by parallel translating Δ with respect to the normal connection of $\widetilde{M}^c$. Set $\widetilde{U} := \bigcup_{u \in U} (\text{Span}\{(n_i)_u \mid i \in I_P \setminus \{0\}\} \cap \Delta_u)$, which is an open subset of the affine subspace $(W_P)_{\gamma(0)}$. Since an element of holonomy along γ is defined on U , there exists the D_P -horizontal curve γ_u parallel to γ with $\gamma_u(0) = u$ for each $u \in U$. Define a map $h_t : \widetilde{U} \rightarrow (W_P)_{\gamma(t)}$ ($0 \leq t \leq 1$) by $h_t(u+w) = \gamma_u(t) + \tau_{\gamma_u|_{[0,t]}}^\perp(w)$ ($u \in U, w \in \text{Span}\{(n_i)_u \mid i \in I_P \setminus \{0\}\} \cap \Delta_u$) (see Fig. 2). It is shown that h_t is a holomorphic isometry into $(W_P)_{\gamma(t)}$ (see Lemma 1.2 in [HL2]). Hence h_t extends to a holomorphic isometry of $(W_P)_{\gamma(0)}$ onto $(W_P)_{\gamma(t)}$. It is shown that this extension is the desired one-parameter family (see Step 3 of the proof of Proposition 1.1 in [HL2]). q.e.d.



Fix $i_0 \in I \cup \{0\}$. Denote by $\Phi_{i_0}(u_0)$ the group of holomorphic isometries of $(W_P)_{u_0}$ generated by

$$\{h_{\gamma,1}^{E_{i_0}} \mid \gamma : E_{i_0}^\perp - \text{horizontal curve s.t. } \gamma(0), \gamma(1) \in L_u^{E_{i_0}}\},$$

where $L_u^{E_{i_0}}$ is the integral manifold of E_{i_0} through u . Also, denote by $\Phi_{i_0}^0(u_0)$ the identity component of $\Phi_{i_0}(u_0)$ and $\Phi_{i_0}^0(u_0)_{u_0}$ the isotropy subgroup of $\Phi_{i_0}^0(u_0)$ at u_0 . Give $\Phi_{i_0}^0(u_0)$ the metric associated with its representation on $(W_P)_{u_0}$. Take $X \in \text{Lie } \Phi_{i_0}^0(u_0) \ominus \text{Lie } \Phi_{i_0}^0(u_0)_{u_0}$, where $\text{Lie}(\cdot)$ is the Lie algebra of $(\cdot)$. Set $g(t) := \exp tX$ and $\gamma(t) := g(t)u_0$, where $\exp$ is the exponential map of $\Phi_{i_0}^0(u_0)$. It is clear that γ is a curve in $L_{u_0}^{E_{i_0}}$. Hence γ is an E_i -horizontal curve for $i \in I$ with $i \neq i_0$. We define a holomorphic isometry F_γ of V by $F_\gamma(\gamma(0)) = \gamma(1)$ and

$$(3.1) \quad (F_\gamma)_{*\gamma(0)} = \begin{cases} g(1)_{*\gamma(0)} & \text{on } (E_{i_0})_{\gamma(0)} \\ (h_{\gamma,1}^{E_i})_{*\gamma(0)} & \text{on } (E_i)_{\gamma(0)} \ (i \neq i_0) \\ \tau_\gamma^\perp & \text{on } T_{\gamma(0)}^\perp \widetilde{M}^c \end{cases}$$

In similar to Theorem 4.1 of [HL2], we have the following fact.

Proposition 3.5. *The holomorphic isometry F_γ preserves $\widetilde{M}^c$ invariantly (i.e., $F_\gamma(\widetilde{M}^c) = \widetilde{M}^c$). Furthermore, it preserves E_i ($i \in I$) invariantly (i.e., $F_{\gamma*}(E_i) = E_i$).*

To show this proposition, we prepare lemmas. By imitating the proof (P163~166) of Proposition 3.1 in [HL2], we can show the following fact.

Lemma 3.5.1. *Let N and $\widehat{N}$ be irreducible proper anti-Kaehlerian isoparametric submanifolds of complex codimension greater than one in an infinite dimensional anti-Kaehlerian space. If $N \cap \widehat{N} \neq \emptyset$ and, for some $p_0 \in N \cap \widehat{N}$, $T_{p_0}N = T_{p_0}\widehat{N}$ and there exists a complex affine line l_0 of $T_{p_0}^\perp N (= T_{p_0}^\perp \widehat{N})$ such that $L_{p_0}^{D_l} = L_{p_0}^{\widehat{D}_l}$ for any complex affine line l of $T_{p_0}^\perp N$ with $l \neq l_0$, then $N = \widehat{N}$ holds, where $L_{p_0}^{D_l}$ (resp. $L_{p_0}^{\widehat{D}_l}$) is the leaf through p_0 of the integrable distribution D_l (resp. $\widehat{D}_l$) on N (resp. $\widehat{N}$) defined as above for l .*

Proof. Let $\{E_i \mid i \in I\} \cup \{E_0\}$ (resp. $\{\widehat{E}_i \mid i \in \widehat{I}\} \cup \{\widehat{E}_0\}$) be the set of all complex curvature distributions of N (resp. $\widehat{N}$). Let $Q_0(p_0)$ (resp. $\widehat{Q}_0(p_0)$) be the set of all points of N (resp. $\widehat{N}$) connected with p_0 by a piecewise smooth curve in N (resp. $\widehat{N}$) each of whose smooth segments is contained in some complex curvature sphere or some integral manifold of E_0 (resp. $\widehat{E}_0$), where E_0 (resp. $\widehat{E}_0$) is the distribution on N (resp. $\widehat{N}$) corresponding to the above distribution E_0 on $\widetilde{M}^c$. Take any $p \in Q_0(p_0)$. There exists a sequence $\{p_0, p_1, \dots, p_l (= p)\}$ such that, for each $j \in \{1, \dots, l\}$, $p_j \in (\bigcup_{i \in I} L_{p_{j-1}}^{E_i}) \cup L_{p_{j-1}}^{E_0}$. Assume

that there exists $j_0 \in \{1, \dots, l\}$ such that $p_{j_0} \in L_{p_{j_0-1}}^{E_{i_0}}$ for some $i_0 \in I$ with $(n_{i_0})_{p_0} \in l_0$. Since N is irreducible (hence the complex Coxeter group associated with N is irreducible) and $\text{codim}_{\mathbb{C}} N \geq 2$, there exists a complex curvature normal n_{i_1} such that $(n_{i_1})_{p_0}$ and $(n_{i_0})_{p_0}$ are linearly independent, $\langle (n_{i_1})_{p_0}, (n_{i_0})_{p_0} \rangle \neq 0$ and that $(n_{i_1})_{p_0}$ does not belong to l_0 . Denote by $l_{i_0 i_1}$ the affine line in $T_{p_0}^\perp N$ through $(n_{i_0})_{p_0}$ and $(n_{i_1})_{p_0}$, and set $D_{i_0 i_1} := D_{l_{i_0 i_1}}$ for simplicity. According to Theorem 3.3, $L_{p_{j_0-1}}^{D_{i_0 i_1}}$ is an irreducible proper anti-Kaehlerian isoparametric submanifold in $(W_{l_{i_0 i_1}})_{p_{j_0-1}}$ of complex codimension two. Hence, by the anti-Kaehlerian version of Theorem D of [HOT], p_{j_0-1} can be joined to p_{j_0} by a piecewise smooth curve each of whose smooth segments is tangent to one of E_i 's ($i \in I$ s.t. $(n_i)_{p_0} \in l_{i_0 i_1}$ and $(n_i)_{p_0} \neq (n_{i_0})_{p_0}$). Therefore, we can take a sequence $\{p_0, p'_1, \dots, p'_{l'} (= p)\}$ such that, for each $j \in \{1, \dots, l'\}$, $p'_j \in \left(\bigcup_{i \in I \text{ s.t. } (n_i)_{p_0} \notin l_0} L_{p'_{j-1}}^{E_i} \right) \cup L_{p'_{j-1}}^{E_0}$. Hence it follows from Lemma 3.5.2 (see below) that $p'_1 \in \widehat{Q}_0(p_0)$, $p'_2 \in \widehat{Q}_0(p'_1)$, $\dots$, $p'_{l'-1} \in \widehat{Q}_0(p'_{l'-2})$ and $p \in \widehat{Q}_0(p'_{l'-1})$ inductively. Therefore we have $p \in \widehat{Q}_0(p_0)$. From the arbitrariness of p , it follows that $Q_0(p_0) \subset \widehat{Q}_0(p_0)$. Similarly we can show $\widehat{Q}_0(p_0) \subset Q_0(p_0)$. Thus we obtain $Q_0(p_0) = \widehat{Q}_0(p_0)$ and hence $\overline{Q_0(p_0)} = \widehat{Q_0(p_0)}$. Let $\mathcal{D}_E^0$ (resp. $\widehat{\mathcal{D}}_E^0$) be the set of all local (smooth) vector fields of N (resp. $\widehat{N}$) which is tangent to some E_i (resp. $\widehat{E}_i$) (where i may be equal to 0) at each point of the domain. For $\mathcal{D}_E^0$ and $p \in N$ (resp. $\widehat{\mathcal{D}}_E^0$ and $\widehat{p} \in \widehat{N}$), define $\Omega_{\mathcal{D}_E^0}(p)$, $(\mathcal{D}_E^0)^*$ and $(\mathcal{D}_E^0)^*(p)$ (resp. $\Omega_{\widehat{\mathcal{D}}_E^0}(\widehat{p})$, $(\widehat{\mathcal{D}}_E^0)^*$ and $(\widehat{\mathcal{D}}_E^0)^*(\widehat{p})$) as the quantities corresponding to the above $\Omega_D(x)$, $\mathcal{D}^*$ and $\mathcal{D}^*(x)$. Since $\overline{(\mathcal{D}_E^0)^*(p)} = \overline{(E_0)_p \oplus (\bigoplus_{i \in I} (E_i)_p)} = T_p N$ for each $p \in N$, it follows from Theorem 3.1 that $\overline{\Omega_{\mathcal{D}_E^0}(p_0)} = N$. Similarly, we have $\overline{\Omega_{\widehat{\mathcal{D}}_E^0}(p_0)} = \widehat{N}$. Also, it is clear that $\Omega_{\mathcal{D}_E^0}(p_0) = Q_0(p_0)$ and $\Omega_{\widehat{\mathcal{D}}_E^0}(p_0) = \widehat{Q}_0(p_0)$. Therefore we obtain $N = \widehat{N}$.

q.e.d.

Lemma 3.5.2. *Let N , $\widehat{N}$, p_0 and l_0 be as in Lemma 3.5.1. Then, for any $p \in L_{p_0}^{E_0} \cup \left(\bigcup_{i \in I \text{ s.t. } (n_i)_{p_0} \notin l_0} L_{p_0}^{E_i} \right)$, we have $T_p N = T_p \widehat{N}$ and $L_p^{D_l} = L_p^{\widehat{D}_l}$ for any complex affine line l of $T_{p_0}^\perp N$ with $l \neq l_0$.*

Proof. First we consider the case where $p \in L_{p_0}^{E_i}$ for some i with $(n_i)_{p_0} \notin l_0$. Then, from the assumption, we have $p \in L_{p_0}^{E_i} = L_{p_0}^{\widehat{E}_i}$ and hence $L_p^{E_i} = L_p^{\widehat{E}_i}$. Let l be a complex affine line of $T_{p_0}^\perp N$ with $l \neq l_0$. Assume that $(n_{i_0})_{p_0} \in l$. Then we have $p \in L_{p_0}^{E_i} \subset L_{p_0}^{D_{l_0}}$. Since $l \neq l_0$, it follows from the assumption that $L_{p_0}^{D_l} = L_{p_0}^{\widehat{D}_l}$. Hence we have $L_p^{D_l} = L_p^{\widehat{D}_l}$. Assume that $(n_i)_{p_0} \notin l$. Take a curve $\gamma : [0, 1] \rightarrow L_{p_0}^{E_i}$ with $\gamma(0) = p_0$ and $\gamma(1) = p$. Since

$(n_i)_{p_0} \notin l$, γ is D_l -horizontal. For the holomorphic isometries $h_{\gamma,1}^{D_l} : (W_l)_{p_0} \rightarrow (W_l)_p$ and $h_{\gamma,1}^{\widehat{D}_l} : (\widehat{W}_l)_{p_0} \rightarrow (\widehat{W}_l)_p$ as in Lemma 3.4, we have $h_{\gamma,1}^{D_l}(L_{p_0}^{D_l}) = L_p^{D_l}$ and $h_{\gamma,1}^{\widehat{D}_l}(L_{p_0}^{\widehat{D}_l}) = L_p^{\widehat{D}_l}$. On the other hand, by imitating the discussion from Line 7 from bottom of Page 164 to Line 4 of Page 165 in [HL2], we can show $h_{\gamma,1}^{D_l} = h_{\gamma,1}^{\widehat{D}_l}$. Hence we obtain $L_p^{D_l} = L_p^{\widehat{D}_l}$. Therefore we have $\sum_{l \neq l_0} T_P L_p^{D_l} = \sum_{l \neq l_0} T_P L_p^{\widehat{D}_l}$. On the other hand, we have $T_p N = \sum_{l \neq l_0} T_P L_p^{D_l}$ and $T_p \widehat{N} = \sum_{l \neq l_0} T_P L_p^{\widehat{D}_l}$. Therefore we have $T_p N = T_p \widehat{N}$. Next we consider the case of $p \in L_{p_0}^{E_0}$. Let l be a complex affine line of $T_{p_0}^\perp N$ with $l \neq l_0$. Assume that $0 \in l$. Then we have $p \in L_{p_0}^{D_l} = L_{p_0}^{\widehat{D}_l}$ and hence $L_p^{D_l} = L_p^{\widehat{D}_l}$. Assume that $0 \notin l$. Take a curve $\gamma : [0, 1] \rightarrow L_{p_0}^{E_0}$ with $\gamma(0) = p_0$ and $\gamma(1) = p$. Since $0 \notin l$, γ is D_l -horizontal. For the holomorphic isometries $h_{\gamma,1}^{D_l} : (W_l)_{p_0} \rightarrow (W_l)_p$ and $h_{\gamma,1}^{\widehat{D}_l} : (\widehat{W}_l)_{p_0} \rightarrow (\widehat{W}_l)_p$, we have $h_{\gamma,1}^{D_l}(L_{p_0}^{D_l}) = L_p^{D_l}$ and $h_{\gamma,1}^{\widehat{D}_l}(L_{p_0}^{\widehat{D}_l}) = L_p^{\widehat{D}_l}$. On the other hand, by imitating the discussion from Line 18 of Page 165 to Line 6 of Page 166 in [HL2], we have $h_{\gamma,1}^{D_l} = h_{\gamma,1}^{\widehat{D}_l}$. Hence we obtain $L_p^{D_l} = L_p^{\widehat{D}_l}$. Therefore we have $\sum_{l \neq l_0} T_P L_p^{D_l} = \sum_{l \neq l_0} T_P L_p^{\widehat{D}_l}$ and hence $T_p N = T_p \widehat{N}$. This completes the proof. q.e.d.

Let N be a proper anti-Kaehlerian isoparametric submanifold in a finite dimensional anti-Kaehlerian space and $\{E_1, \dots, E_k\}$ be the set of all complex curvature distributions of N . We can define the holomorphic isometry of the anti-Kaehlerian space corresponding to the above F_γ . Denote by the same symbol F_γ this holomorphic isometry. In similar to Lemma 4.2 in [HL2], we have the following fact.

Lemma 3.5.3. *If N is a principal orbit through a semi-simple element of an aks-representation, then N is proper anti-Kaehlerian isoparametric and $F_\gamma(N) = N$ holds.*

Proof. First we note that any irreducible (semi-simple) anti-Kaehlerian symmetric space is regarded as the complexification of an irreducible Riemannian symmetric space of non-compact type. Let L/H be an irreducible Riemannian symmetric space of non-compact type. Denote by $\mathfrak{l}$ (resp. $\mathfrak{h}$) the Lie algebra of L (resp. H). Let θ be the Cartan involution of L with $(\text{Fix } \theta)_0 \subset H \subset \text{Fix } \theta$ and denote by the same symbol θ the involution of $\mathfrak{l}$ associated with θ . Set $\mathfrak{q} := \text{Ker}(\theta + \text{id})$, which is identified with the tangent space $T_{eH}(L/H)$. The complexification $\mathfrak{q}^\mathbb{C}$ is identified with the tangent space $T_{eH^\mathbb{C}}(L^\mathbb{C}/H^\mathbb{C})$ of the associated anti-Kaehlerian symmetric space $L^\mathbb{C}/H^\mathbb{C}$. Let ρ be the aks-representation associated with $L^\mathbb{C}/H^\mathbb{C}$ and N be a principal orbit of the representation ρ through a semi-simple element $w \in \mathfrak{q}^\mathbb{C}$, that is, $N = \rho(H^\mathbb{C}) \cdot w$. Denote by A the shape tensor of N . Let $\mathfrak{a}$ be a Cartan subspace of $\mathfrak{q}^\mathbb{C}$ containing w . The space $\mathfrak{a}$ contains the maximal split abelian

subspace $\mathfrak{a}_v := \mathfrak{a} \cap \mathfrak{q}$ of vector-type and $\mathfrak{a} = \mathfrak{a}_v^{\mathbf{c}}$ holds. For each (**R**-)linear function α on $\mathfrak{a}_v$ (i.e., $\alpha \in \mathfrak{a}_v^*$), we set

$$\mathfrak{q}_\alpha^{\mathbf{c}} := \{X \in \mathfrak{q}^{\mathbf{c}} \mid \text{ad}(a)^2(X) = \alpha(a)^2 X \ (\forall a \in \mathfrak{a}_v)\}.$$

Set $\Delta := \{\alpha \in \mathfrak{a}_v^* \mid \mathfrak{q}_\alpha^{\mathbf{c}} \neq \{0\}\}$, which is called the root system with respect to $\mathfrak{a}_v$. Then we have the root space decomposition

$$\mathfrak{q}^{\mathbf{c}} = \mathfrak{a} + \sum_{\alpha \in \Delta_+} \mathfrak{q}_\alpha^{\mathbf{c}},$$

where Δ_+ is the positive root system under some lexicographical ordering of Δ and we note that $\mathfrak{a}$ is equal to the centralizer of $\mathfrak{a}_v$ in $\mathfrak{q}^{\mathbf{c}}$. For each $\alpha \in \Delta_+$, the complexification $\alpha^{\mathbf{c}}$ is regarded as a **C**-linear function on $\mathfrak{a}$ and we have

$$\mathfrak{q}_\alpha^{\mathbf{c}} = \{X \in \mathfrak{q}^{\mathbf{c}} \mid \text{ad}(a)^2(X) = \alpha^{\mathbf{c}}(a)^2 X \ (\forall a \in \mathfrak{a})\}.$$

Since N is a principal orbit and hence w is a regular element, we have $\alpha^{\mathbf{c}}(w) \neq 0$ for any $\alpha \in \Delta_+$. Under the identification of $\mathfrak{a}$ with $T_w^\perp N$, $\alpha^{\mathbf{c}}$ is regarded as a **C**-linear function on $T_w^\perp N$, which is denoted by $\overline{\alpha^{\mathbf{c}}}$. Easily we can show

$$A_v^{\mathbf{c}}|_{\mathfrak{q}_\alpha^{\mathbf{c}}} = -\frac{\overline{\alpha^{\mathbf{c}}}(v)}{\alpha^{\mathbf{c}}(w)} \text{id} \quad (\alpha \in \Delta_+)$$

for any $v \in T_w^\perp N$. Let $\lambda_{\alpha^{\mathbf{c}}, w}$ be the parallel section of the **C**-dual bundle $(T^\perp N)^*$ of $T^\perp N$ with $(\lambda_{\alpha^{\mathbf{c}}, w})_w = -\frac{1}{\alpha^{\mathbf{c}}(w)} \overline{\alpha^{\mathbf{c}}}$. It is clear that N is a proper anti-Kaehlerian isoparametric submanifold having $\{\lambda_{\alpha^{\mathbf{c}}, w} \mid \alpha \in \Delta_+\}$ as the set of all complex principal curvatures. Denote by $E_{\alpha^{\mathbf{c}}}$ the complex curvature distribution for $\lambda_{\alpha^{\mathbf{c}}, w}$. Take $v_0 \in (\lambda_{\alpha_0^{\mathbf{c}}, w})^{-1}(1) \setminus (\bigcup_{\substack{\alpha \in \Delta_+ \\ \text{s.t. } \alpha \neq \alpha_0}} (\lambda_{\alpha^{\mathbf{c}}, w})^{-1}(1))$ and set $F := \rho(H^{\mathbf{c}}) \cdot v_0$, which is a focal submanifold of N whose corresponding focal distribution is equal to $E_{\alpha_0^{\mathbf{c}}}$. We have the relations $\mathfrak{h}^{\mathbf{c}} = \mathfrak{z}_{\mathfrak{h}^{\mathbf{c}}}(\mathfrak{a}_v) + \sum_{\alpha \in \Delta_+} \mathfrak{h}_\alpha^{\mathbf{c}}$, $T_{v_0} F = \sum_{\substack{\alpha \in \Delta_+ \\ \text{s.t. } \alpha \neq \alpha_0}} \mathfrak{q}_\alpha^{\mathbf{c}}$ and $T_{v_0}^\perp F = \mathfrak{a} + \mathfrak{q}_{\alpha_0}^{\mathbf{c}}$, where $\mathfrak{z}_{\mathfrak{h}^{\mathbf{c}}}(\mathfrak{a}_v)$

is the centralizer of $\mathfrak{a}_v$ in $\mathfrak{h}^{\mathbf{c}}$ and $\mathfrak{h}_\alpha^{\mathbf{c}} := \{X \in \mathfrak{h}^{\mathbf{c}} \mid \text{ad}(a)^2 X = \alpha(a)^2 X \ (\forall a \in \mathfrak{a}_v)\}$. Denote by $H_w^{\mathbf{c}}$ (resp. $H_{v_0}^{\mathbf{c}}$) the isotropy group of the $H^{\mathbf{c}}$ -action at w (resp. v_0) and by $\mathfrak{h}_w^{\mathbf{c}}$ (resp. $\mathfrak{h}_{v_0}^{\mathbf{c}}$) the Lie algebra of $H_w^{\mathbf{c}}$ (resp. $H_{v_0}^{\mathbf{c}}$). Then we have $\mathfrak{h}_w^{\mathbf{c}} = \mathfrak{z}_{\mathfrak{h}^{\mathbf{c}}}(\mathfrak{a}_v)$ and $\mathfrak{h}_{v_0}^{\mathbf{c}} = \mathfrak{z}_{\mathfrak{h}^{\mathbf{c}}}(\mathfrak{a}_v) + \mathfrak{h}_{\alpha_0}^{\mathbf{c}}$. For the restriction of the $\rho(H_{v_0}^{\mathbf{c}})$ -action on $\mathfrak{q}^{\mathbf{c}}$ to $T_{v_0}^\perp F$ is called the slice representation of the action at v_0 . It is shown that this slice representation coincides with the normal holonomy group action of F at v_0 and $\rho(H_{v_0}^{\mathbf{c}}) \cdot w = L_w^{E_{\alpha_0^{\mathbf{c}}}}$. Set $\Phi(v_0) := \rho(H_{v_0}^{\mathbf{c}})$ and $\Phi(w) := \rho(H_w^{\mathbf{c}})$.

The leaf $L_w^{E_{\alpha_0^{\mathbf{c}}}}$ is regarded as the quotient manifold $\Phi(v_0)/\Phi(w)$. The holomorphic isometry F_γ in the statement is given as follows. Take $X (= \text{ad}_{\mathfrak{r}}(\overline{X})) \in \text{Lie } \Phi(v_0) \ominus \text{Lie } \Phi(w)$, where $\overline{X} \in \mathfrak{h}_{v_0}^{\mathbf{c}}$, and set $g(t) := \exp_{\Phi(v_0)}(tX)$ and $\gamma(t) := g(t) \cdot w$, where $t \in [0, 1]$. Then

F_γ is given by $F_\gamma(w) = \gamma(1)$, $F_{\gamma*}w|_{(E_{\alpha_0^c})_w} = g(1)*w|_{(E_{\alpha_0^c})_w}$, $F_{\gamma*}w|_{(E_{\alpha^c})_w} = h_{\gamma,1}^{E_{\alpha^c}}|_{(E_{\alpha^c})_w}$ ($\alpha \in \Delta_+$ s.t. $\alpha \neq \alpha_0$) and $F_{\gamma*}w|_{T_w^\perp N} = \tau_\gamma^\perp$, where $h_{\gamma,1}^{E_{\alpha^c}}$ is defined as in Lemm 3.4 and $\tau_\gamma^\perp$ is the parallel translation along γ with respect to the normal connection of N . Easily we can show $(F_\gamma)*w = g(1)*w$. Hence, since both F_γ and $g(1)$ are affine transformations of $\mathfrak{q}^c$, they coincide with each other. Therefore, we obtain $F_\gamma(N) = g(1)(\rho(H^c) \cdot w) = \rho(\exp_{L^c}(\bar{X}))(\rho(H^c) \cdot w) = N$.

q.e.d.

By using Lemmas 3.5.1 and 3.5.3, we shall prove Proposition 3.5.

Proof of Proposition 3.5. Since M is an irreducible proper complex equifocal submanifold, $\widetilde{M}^c$ is a full irreducible proper anti-Kaehlerian isoparametric submanifold. Since F_γ is an isometry, so is also $\widetilde{M}^{c'} := F_\gamma(\widetilde{M}^c)$. Let $\{E'_i | i \in I\} \cup \{E'_0\}$ be the set of all complex curvature distributions on $\widetilde{M}^{c'}$ and n'_i be the complex curvature normal corresponding to E'_i . From the definition of F_γ , it follows that $\gamma(1) \in \widetilde{M}^c \cap \widetilde{M}^{c'}$, $T_{\gamma(1)}\widetilde{M}^c = T_{\gamma(1)}\widetilde{M}^{c'}$, complex curvature normals of $\widetilde{M}^{c'}$ coincide with those of $\widetilde{M}^c$ at $\gamma(1)$ and that complex curvature spheres of $\widetilde{M}^{c'}$ through $\gamma(1)$ coincides with those of $\widetilde{M}^c$ through $\gamma(1)$. Let l_0 be the complex affine line through 0 and $(n_{i_0})_{\gamma(1)}$, that is, $l_0 := \text{Span}_{\mathbf{C}}\{(n_{i_0})_{\gamma(1)}\}$. Let l be any complex affine line of $T_{\gamma(1)}^\perp \widetilde{M}^c$ with $l \neq l_0$. Now we shall show that $L_{\gamma(1)}^{D_l} = L_{\gamma(1)}^{D'_l}$, where D_l (resp. D'_l) is the distribution on $\widetilde{M}^c$ (resp. $\widetilde{M}^{c'}$) defined as above for l . First we consider the case of $(n_{i_0})_{\gamma(1)} \in l$. Then we have $0 \notin l$. If there does not exist $i_1 (\neq i_0) \in I$ with $(n_{i_1})_{\gamma(1)} \in l$, then we have $L_{\gamma(1)}^{D_l} = L_{\gamma(1)}^{E_{i_0}} = L_{\gamma(1)}^{E'_{i_0}} = L_{\gamma(1)}^{D'_l}$. Assume that there exists $i_1 (\neq i_0) \in I$ with $(n_{i_1})_{\gamma(1)} \in l$. Let v be a complex focal normal vector field of $\widetilde{M}^c$ such that the corresponding focal distribution is equal to D_l . Since $0 \notin l$, it follows from Theorem 3.3 that $L_{\gamma(1)}^{D_l}$ is a principal orbit of an aks-representation on $T_{f_v(\gamma(1))}^\perp F_v$. Since $(n_{i_0})_{\gamma(1)}, (n_{i_1})_{\gamma(1)} \in l$ and $0 \notin l$, $(n_{i_0})_{\gamma(1)}$ and $(n_{i_1})_{\gamma(1)}$ are $\mathbf{C}$ -linear independent. Hence the aks-representation is the one associated with some anti-Kaehlerian symmetric space of complex rank two. If $L_{\gamma(1)}^{D_l}$ is reducible, then the complex Coxeter group associated with $L_{\gamma(1)}^{D_l} (\subset T_{f_v(\gamma(1))}^\perp F_v)$ is reducible. Hence it follows from Lemma 3.8 of [Koi4] that $(n_{i_0})_{\gamma(1)}$ and $(n_{i_1})_{\gamma(1)}$ are orthogonal and that, for any complex curvature normal n of $L_{\gamma(1)}^{D_l}$, $n_{\gamma(1)}$ is contained in $l_0 \cup \text{Span}_{\mathbf{C}}\{(n_{i_1})_{\gamma(1)}\}$. Also, since $n_{\gamma(1)} \in l$, $n_{\gamma(1)}$ is equal to $(n_{i_0})_{\gamma(1)}$ or $(n_{i_1})_{\gamma(1)}$. Hence the set of all complex curvature normals of $L_{\gamma(1)}^{D_l}$ is equal to $\{n_{i_0}|_{L_{\gamma(1)}^{D_l}}, n_{i_1}|_{L_{\gamma(1)}^{D_l}}\}$. This implies that $L_{\gamma(1)}^{D_l}$ is congruent to the (extrinsic) product of complex spheres $L_{\gamma(1)}^{E_{i_0}}$ and $L_{\gamma(1)}^{E_{i_1}}$. Similarly $L_{u_0}^{D_l}$ is congruent to the (extrinsic) product of

$L_{u_0}^{E_{i_0}}$ and $L_{u_0}^{E_{i_1}}$. Hence we have

$$F_\gamma(L_{u_0}^{D_l}) = F_\gamma(L_{u_0}^{E_{i_0}}) \times F_\gamma(L_{u_0}^{E_{i_1}}) = L_{\gamma(1)}^{E_{i_0}} \times L_{\gamma(1)}^{E_{i_1}} = L_{\gamma(1)}^{D_l}.$$

On the other hand, we have $F_\gamma(L_{u_0}^{D_l}) = L_{\gamma(1)}^{D'_l}$. Therefore we obtain $L_{\gamma(1)}^{D_l} = L_{\gamma(1)}^{D'_l}$. If $L_{\gamma(1)}^{D_l}$ is irreducible, then it follows from Lemma 3.5.3 that $F_\gamma(L_{\gamma(1)}^{D_l}) = (F_\gamma|_{(W_l)_{\gamma(1)}})(L_{\gamma(1)}^{D_l}) = L_{\gamma(1)}^{D_l}$. Hence we obtain $L_{\gamma(1)}^{D_l} = L_{\gamma(1)}^{D'_l}$. Next we consider the case of $(n_{i_0})_{\gamma(1)} \notin l$. Then, since γ is D_l -horizontal, we have $F_\gamma(L_{u_0}^{D_l}) = L_{\gamma(1)}^{D_l}$ and hence $L_{\gamma(1)}^{D_l} = L_{\gamma(1)}^{D'_l}$. Therefore, from Lemma 3.5.1, we obtain $F_\gamma(\widetilde{M}^c) = \widetilde{M}^c$. Furthermore, from this relation, we can show $(F_\gamma)_*(E_i) = E_i$ ($i \in I$) easily. q.e.d.

By using Proposition 3.5, we prove the following fact.

Proposition 3.6. *For any $u \in Q(u_0)$, there exists a holomorphic isometry f of V such that $f(u_0) = u$, $f(\widetilde{M}^c) = \widetilde{M}^c$ (hence $f_*(E_i) = E_i$ ($i \in I$)), $f(Q(u_0)) = Q(u_0)$ and that $f_{*u_0}|_{T_{u_0}^\perp \widetilde{M}^c}$ coincides with the parallel translation along a curve in $\widetilde{M}^c$ starting from u_0 and terminating to u with respect to the normal connection of $\widetilde{M}^c$.*

Proof. Take a sequence $\{u_0, u_1, \dots, u_k (= u)\}$ of $Q(u_0)$ such that, for each $i \in \{0, 1, \dots, k-1\}$, u_i and u_{i+1} are some complex curvature sphere S_i^c of $\widetilde{M}^c$. Furthermore, for each $i \in \{0, 1, \dots, k-1\}$, we take the geodesic $\gamma_i : [0, 1] \rightarrow S_i^c$ with $\gamma_i(0) = u_i$ and $\gamma_i(1) = u_{i+1}$, and define a holomorphic isometry F_{γ_i} of V as in (3.1). Set $f := F_{\gamma_{k-1}} \circ \dots \circ F_{\gamma_1} \circ F_{\gamma_0}$. According to the definition of F_{γ_i} ($i = 0, 1, \dots, k-1$) and Proposition 3.5, f preserves $\widetilde{M}^c$ invariantly and the restriction of f_{*u_0} to $T_{u_0}^\perp \widetilde{M}^c$ coincides with the parallel translation with respect to the normal connection of $\widetilde{M}^c$. Also, since f preserves complex curvature spheres invariantly, it is shown that f preserves $Q(u_0)$ invariantly. Thus f is the desired holomorphic isometry. q.e.d.

By using Propositions 3.2 and 3.6, we shall prove the homogeneity of $\widetilde{M}^c$.

Proposition 3.7. *$\widetilde{M}^c$ is extrinsically homogeneous.*

Proof. Take any $\widehat{u} \in \widetilde{M}^c$. Since $\overline{Q(u_0)} = \widetilde{M}^c$ by Proposition 3.2, there exists a sequence $\{u_k\}_{k=1}^\infty$ in $Q(u_0)$ with $\lim_{k \rightarrow \infty} u_k = \widehat{u}$. According to Proposition 3.6, for each $k \in \mathbf{N}$, there

exists a holomorphic isometry f_k of V with $f_k(u_0) = u_k$, $f_k(\widetilde{M^c}) = \widetilde{M^c}$, $f_k(Q(u_0)) = Q(u_0)$ and $f_k(L_{u_0}^{E_i}) = L_{u_k}^{E_i}$ ($i \in I$).

(Step I) First we shall show that, for each $i \in I$, there exists a subsequence $\{f_{k_j}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ such that $\{f_{k_j}|_{L_{u_0}^{E_i}}\}_{j=1}^\infty$ pointwisely converges to some holomorphic isometry of $L_{u_0}^{E_i}$ onto $L_{\widehat{u}}^{E_i}$. Let $(L_{u_0}^{E_i})_{\mathbf{R}}$ be the compact real of the complex sphere $L_{u_0}^{E_i}$ satisfying $\langle T_{u_0}(L_{u_0}^{E_i})_{\mathbf{R}}, JT_{u_0}(L_{u_0}^{E_i})_{\mathbf{R}} \rangle = 0$. Also, let $(L_{u_k}^{E_i})_{\mathbf{R}}$ and $(L_{\widehat{u}}^{E_i})_{\mathbf{R}}$ be the same kind of compact reals of $L_{u_k}^{E_i}$ and $L_{\widehat{u}}^{E_i}$, respectively. Clearly we have $f_k((L_{u_0}^{E_i})_{\mathbf{R}}) = (L_{u_k}^{E_i})_{\mathbf{R}}$. Let $\widetilde{M^c}/\mathfrak{F}_i$ be the leaf space of the foliation $\mathfrak{F}_i$ consisting of the integral manifolds of E_i and $\psi_i : \widetilde{M^c} \rightarrow \widetilde{M^c}/\mathfrak{F}_i$ be the quotient map. Take a sufficiently small tubular neighborhood U of $L_{\widehat{u}}^{E_i}$ in V such that $L_u^{E_i} \subset U$ for each $u \in U$. Let $m_i := \dim E_i$. Take a base $\{e_1, \dots, e_{m_i}\}$ of $T_{u_0}((L_{u_0}^{E_i})_{\mathbf{R}})$ such that the norms $\|e_1\|, \dots, \|e_{m_i}\|$ are sufficiently small and $\bar{u}_a := \exp(e_a)$ ($a = 1, \dots, m_i$), where $\exp$ is the exponential map of $(L_{u_0}^{E_i})_{\mathbf{R}}$. Since $\lim_{k \rightarrow \infty} f_k(u_0) = \widehat{u}$ and $(L_u^{E_i})_{\mathbf{R}}$'s ($u \in U$) are compact, there exists a subsequence $\{f_{k_j}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ such that $\{f_{k_j}(\bar{u}_a)\}_{j=1}^\infty$ ($a = 1, \dots, m_i$) converge. Set $\widehat{u}_a := \lim_{j \rightarrow \infty} f_{k_j}(\bar{u}_a)$ ($a = 1, \dots, m_i$). Since $\lim_{j \rightarrow \infty} f_{k_j}(u_0) = \widehat{u}$ and $f_{k_j}((L_{u_0}^{E_i})_{\mathbf{R}}) = (L_{u_{k_j}}^{E_i})_{\mathbf{R}}$, we have $\widehat{u}_a \in (L_{\widehat{u}}^{E_i})_{\mathbf{R}}$ ($a = 1, \dots, m_i$). Denote by d_0, d_j ($j \in \mathbf{N}$) and $\widehat{d}$ the (Riemannian) distance functions of $(L_{u_0}^{E_i})_{\mathbf{R}}$, $(L_{u_{k_j}}^{E_i})_{\mathbf{R}}$ and $(L_{\widehat{u}}^{E_i})_{\mathbf{R}}$, respectively. Since each $f_{k_j}|_{(L_{u_0}^{E_i})_{\mathbf{R}}}$ is an isometry onto $(L_{u_{k_j}}^{E_i})_{\mathbf{R}}$, we have $d_j(f_{k_j}(u_0), f_{k_j}(\bar{u}_a)) = d_0(u_0, \bar{u}_a)$ and $d_j(f_{k_j}(\bar{u}_a), f_{k_j}(\bar{u}_b)) = d_0(\bar{u}_a, \bar{u}_b)$, ($a, b = 1, \dots, m_i$). Hence we have $\widehat{d}(\widehat{u}, \widehat{u}_a) = d_0(u_0, \bar{u}_a)$ and $\widehat{d}(\widehat{u}_a, \widehat{u}_b) = d_0(\bar{u}_a, \bar{u}_b)$ ($a, b = 1, \dots, m_i$). Therefore there exists a unique holomorphic isometry $\bar{f}$ of $(L_{u_0}^{E_i})_{\mathbf{R}}$ onto $(L_{\widehat{u}}^{E_i})_{\mathbf{R}}$ satisfying $\bar{f}(u_0) = \widehat{u}$ and $\bar{f}(\bar{u}_a) = \widehat{u}_a$ ($a = 1, \dots, m_i$). It is clear that $\bar{f}$ is uniquely extended to a holomorphic isometry of $L_{u_0}^{E_i}$ onto $L_{\widehat{u}}^{E_i}$. Denote by f this holomorphic extension. It is easy to show that $\{f_{k_j}|_{L_{u_0}^{E_i}}\}_{j=1}^\infty$ pointwisely converges to f .

(Step II) Next we shall show that, for each fixed $w \in Q(u_0)$, there exists a subsequence $\{f_{k_j}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ such that $\lim_{j \rightarrow \infty} f_{k_j}(w)$ exists. There exists a sequence $\{u_1, \dots, u_m (= w)\}$ in $Q(u_0)$ such that, for each $j \in \{1, \dots, m\}$, u_j is contained in some complex curvature sphere $L_{u_{j-1}}^{E_{i(j)}}$. For simplicity, assume that $m = 3$. From the fact in Step I, there exists a subsequence $\{f_{k_j^1}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ (in Step I) such that $\{f_{k_j^1}|_{L_{u_0}^{E_{i(1)}}}\}_{j=1}^\infty$ pointwisely converges to some holomorphic isometry f^1 of $L_{u_0}^{E_{i(1)}}$ onto $L_{\widehat{u}}^{E_{i(1)}}$. Again, from the fact in Step I, there exists a subsequence $\{f_{k_j^2}\}_{j=1}^\infty$ of $\{f_{k_j^1}\}_{j=1}^\infty$ such that $\{f_{k_j^2}|_{L_{u_1}^{E_{i(2)}}}\}_{j=1}^\infty$ pointwisely converges to some holomorphic isometry f^2 of $L_{u_1}^{E_{i(2)}}$ onto $L_{f^1(u_1)}^{E_{i(2)}}$. Again, from the fact in step I, there exists a subsequence $\{f_{k_j^3}\}_{j=1}^\infty$ of $\{f_{k_j^2}\}_{j=1}^\infty$ such that $\{f_{k_j^3}|_{L_{u_2}^{E_{i(3)}}}\}_{j=1}^\infty$ pointwisely converges to some holomorphic isometry f^3 of $L_{u_2}^{E_{i(3)}}$ onto $L_{f^2(u_2)}^{E_{i(3)}}$. In particular, we have

$$\lim_{j \rightarrow \infty} f_{k_j^3}(w) = f^3(w).$$

(Step III) Let W be the affine span of $\widetilde{M}^c$. Next we shall show that there exists a subsequence $\{f_{k_j}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ such that $\{f_{k_j}|_W\}_{j=1}^\infty$ pointwisely converges to some holomorphic isometry of W . Take a countable subset $B := \{w_j \mid j \in \mathbf{N}\}$ of $Q(u_0)$ with $\overline{B} = \overline{Q(u_0)} (= \widetilde{M}^c)$. According to the fact in Step II, there exists a subsequence $\{f_{k_j^1}\}_{j=1}^\infty$ of $\{f_k\}_{k=1}^\infty$ such that $\lim_{j \rightarrow \infty} f_{k_j^1}(w_1)$ exists. Again, according to the fact in Step II, there exists a subsequence $\{f_{k_j^2}\}_{j=1}^\infty$ of $\{f_{k_j^1}\}_{j=1}^\infty$ such that $\lim_{j \rightarrow \infty} f_{k_j^2}(w_2)$ exists. In the sequel, we take subsequences $\{f_{k_j^l}\}_{j=1}^\infty$ ($l = 3, 4, 5, \dots$) inductively. It is clear that $\lim_{j \rightarrow \infty} f_{k_j^j}(w_l)$ exists for each $l \in \mathbf{N}$, that is, $\{f_{k_j^j}|_B\}_{j=1}^\infty$ pointwisely converges to some map f of B into $\widetilde{M}^c$. Since each $f_{k_j^j}$ is a holomorphic isometry, f is extended to a holomorphic isometry of $\widetilde{M}^c$. Denote by $\widetilde{f}$ this extension. It is clear that $\{f_{k_j^j}|_{\widetilde{M}^c}\}_{j=1}^\infty$ pointwisely converges to $\widetilde{f}$. Furthermore, since each $f_{k_j^j}$ is an affine transformation and hence the restriction $f_{k_j^j}|_W$ of $f_{k_j^j}$ to W is a holomorphic isometry of W , $\widetilde{f}$ is extended to a holomorphic isometry of W . Denote by $\widetilde{\widetilde{f}}$ this holomorphic extension. It is clear that $\{f_{k_j^j}|_W\}_{j=1}^\infty$ pointwisely converges to $\widetilde{\widetilde{f}}$.

(Step IV) Let f , $\widetilde{f}$ and $\widetilde{\widetilde{f}}$ be as in Step III. It is clear that $\widetilde{\widetilde{f}}$ is extended to a holomorphic isometry of V . Denote by $\widehat{f}$ this extension. We have

$$\widehat{f}(\widetilde{M}^c) = \widetilde{f}(\widetilde{M}^c) = \widetilde{M}^c.$$

Also we have

$$\widehat{f}(u_0) = f(u_0) = \lim_{j \rightarrow \infty} f_{k_j^j}(u_0) = \lim_{j \rightarrow \infty} u_{k_j^j} = \widehat{u}.$$

Let H be the closed subgroup of $I_h(V)$ generated by all holomorphic isometries of V preserving $\widetilde{M}^c$ invariantly, where $I_h(V)$ is the holomorphic isometry group of V . Then it follows from the above fact (together with the arbitrariness of $\widehat{u}$) that $H \cdot u_0 = \widetilde{M}^c$.

q.e.d.

4 Proofs of Theorem A and Corollary B

In this section, we shall prove Theorem A and Corollary B. Let M be an irreducible proper complex equifocal submanifold in a symmetric space G/K of non-compact type. Assume that the codimension of M is greater than one. Let M^c be the (complete extrinsic) complexification of M , $\widehat{M}^c := \pi^c{}^{-1}(M^c)$ and $\widetilde{M}^c := (\pi^c \circ \phi^c)^{-1}(M^c)$, where π^c is the natural

projection of $G^{\mathbf{c}}$ onto $G^{\mathbf{c}}/K^{\mathbf{c}}$ and $\phi^{\mathbf{c}}$ is the parallel transport map for $G^{\mathbf{c}}$. Without loss of generality, we may assume that $K^{\mathbf{c}}$ is connected and that $G^{\mathbf{c}}$ is simply connected. Hence both $\widehat{M}^{\mathbf{c}}$ and $\widetilde{M}^{\mathbf{c}}$ are connected. For simplicity, set $V := H^0([0, 1], \mathfrak{g}^{\mathbf{c}})$. According to Proposition 3.7, $\widetilde{M}^{\mathbf{c}}$ is extrinsically homogeneous, that is, there exists a closed connected subgroup H of the anti-Kaehlerian transformation group (i.e., the holomorphic isometry group) $I_h(V)$ of V having $\widetilde{M}^{\mathbf{c}}$ as an orbit. Let $\rho : H^1([0, 1], G^{\mathbf{c}}) \rightarrow I_h(V)$ be the representation of $H^1([0, 1], G^{\mathbf{c}})$ defined by $\rho(g) := g * \cdot$ ($g \in H^1([0, 1], G^{\mathbf{c}})$). In the proof of Theorem A, it is key to show the following fact.

Proposition 4.1. *The above group H is a subgroup of $\rho(H^1([0, 1], G^{\mathbf{c}}))$.*

To prove this proposition, we prepare some lemmas. Let $\mathcal{K}^h$ be the Lie algebra of all holomorphic Killing fields on V (i.e., the Lie algebra of $I_h(V)$) and $\mathcal{K}_{\widetilde{M}^{\mathbf{c}}}^h$ be the Lie algebra of all holomorphic Killing fields on V which are tangent to $\widetilde{M}^{\mathbf{c}}$ along $\widetilde{M}^{\mathbf{c}}$. For $K \in \mathcal{K}^h$, we define a map $F_K : \Omega_e(G^{\mathbf{c}}) \rightarrow \mathfrak{g}^{\mathbf{c}}$ by $F_K(g) := \phi_{*\hat{0}}^{\mathbf{c}}(((g * \cdot)_* K)_{\hat{0}})$. For this map F_K , we have the following fact.

Lemma 4.1.1. (i) *For $g \in \Omega_e(G^{\mathbf{c}})$, $F_K(g) = \int_0^1 \text{Ad}(g)(K_{g^{-1}*\hat{0}})dt$.*
(ii) *If $K \in \mathcal{K}_{\widetilde{M}^{\mathbf{c}}}^h$, then the image of F_K is contained in $T_e\widehat{M}^{\mathbf{c}}$.*

Proof. Let $\{\psi_s\}_{s \in \mathbf{R}}$ be the one-parameter transformation group associated with K . For each $g \in \Omega_e(G^{\mathbf{c}})$, we have

$$\begin{aligned} ((g * \cdot)_* K)_{\hat{0}} &= \frac{d}{ds} \Big|_{s=0} g * (\psi_s(g^{-1} * \hat{0})) \\ &= \frac{d}{ds} \Big|_{s=0} (\text{Ad}(g)(\psi_s(g^{-1} * \hat{0})) - g'g_*^{-1}) = \text{Ad}(g)(K_{g^{-1}*\hat{0}}). \end{aligned}$$

On the other hand, we have $\phi_{*\hat{0}}^{\mathbf{c}}(u) = \int_0^1 u(t)dt$ ($u \in T_{\hat{0}}V (= V)$) (see Lemma 6 of [Koi2]). Hence we obtain the relation in (i). Since $g \in \Omega_e(G^{\mathbf{c}})$, it maps each fibre of $\phi^{\mathbf{c}}$ to oneself. Hence, if $K \in \mathcal{K}_{\widetilde{M}^{\mathbf{c}}}^h$, then $(g * \cdot)_* K \in \mathcal{K}_{\widetilde{M}^{\mathbf{c}}}^h$. In particular, we have $((g * \cdot)_* K)_{\hat{0}} \in T_{\hat{0}}\widetilde{M}^{\mathbf{c}}$. Therefore we obtain $F_K(g) \in \phi_{*\hat{0}}(T_{\hat{0}}\widetilde{M}^{\mathbf{c}}) = T_e\widehat{M}^{\mathbf{c}}$. q.e.d.

For $v \in H^1([0, 1], \mathfrak{g}^{\mathbf{c}})$, we define a vector field K^v on V by $(K^v)_u := [v, u] - v'$ ($u \in V$). Let $\{\exp sv \mid s \in \mathbf{R}\}$ be the one-parameter subgroup of $H^1([0, 1], G^{\mathbf{c}})$ associated with v . Then the holomorphic Killing field associated with the one-parameter transformation group $\{\rho(\exp sv) \mid s \in \mathbf{R}\}$ of V is equal to K^v . Thus we have $K^v \in \mathcal{K}^h$. For K^v , we have the following fact.

Lemma 4.1.2. *The map F_{K^v} is a constant map.*

Proof. Since $\rho(\exp sv)$ maps the fibres of ϕ^c to them, for $u_1, u_2 \in V$ with $\phi^c(u_1) = \phi^c(u_2)$, we have $\phi^c(\rho(\exp sv)(u_1)) = \phi^c(\rho(\exp sv)(u_2))$ and hence $\phi_{*u_1}^c((K^v)_{u_1}) = \phi_{*u_2}^c((K^v)_{u_2})$. Take $g_1, g_2 \in \Omega_e(G^c)$. Since $g_i * \cdot$ maps each fibre of ϕ^c to oneself, we have $F_{K^v}(g_i) = \phi_{*\hat{0}}^c((\rho(g_i)_*(K^v))_{\hat{0}}) = \phi_{*(g_i^{-1}*\hat{0})}^c((K^v)_{g_i^{-1}*\hat{0}})$. Also, we have $\phi^c(g_1^{-1}*\hat{0}) = \phi^c(g_2^{-1}*\hat{0}) (= e)$. Hence we have $\phi_{*(g_1*\hat{0})}^c((K^v)_{g_1^{-1}*\hat{0}}) = \phi_{*(g_2^{-1}*\hat{0})}^c((K^v)_{g_2^{-1}*\hat{0}})$. Therefore we obtain $F_{K^v}(g_1) = F_{K^v}(g_2)$. Thus F_{K^v} is a constant map. q.e.d.

Also we have the following fact for F_K .

Lemma 4.1.3. (i) *The map $K \mapsto F_K$ is linear.*

(ii) $F_K(g_1 g_2) = F_{\rho(g_2)_* K}(g_1)$ ($g_1, g_2 \in \Omega_e(G^c)$).

(iii) $(dF_K)_g \circ (dR_g)_{\hat{e}} = (dF_{\rho(g)_* K})_{\hat{e}}$ ($g \in \Omega_e(G^c)$).

(iv) *If $K_u = Au + b$ ($u \in V$) for some linear transformation A of V and some $b \in V$, then $(dF_K)_{\hat{e}}(u) = \int_0^1 (A + \text{ad}(\tilde{b}))u' dt$, where $\tilde{b}(t) := \int_0^t b(t) dt$.*

(v) *If $\overline{K} = K + K^v$, then $F_{\overline{K}} - F_K$ is a constant map.*

Proof. The statements (i) $\sim$ (iii) are trivial. The statement (iv) is shown by imitating the proof of Proposition 2.3 of [Ch]. For $u \in T_{\hat{e}}(\Omega_e(G^c)) \subset H^1([0, 1], \mathfrak{g}^c)$, it follows from (iv) that

$$d(F_{\overline{K}} - F_K)_{\hat{e}}(u) = (dF_{K^v})_{\hat{e}}(u) = \int_0^1 (\text{ad}(v) + \text{ad}(-v))u' dt = 0.$$

This implies that $F_{\overline{K}} - F_K$ is a constant map. Thus the statement (v) follows. q.e.d.

By imitating the proof of Theorem 2.2 of [Ch], we can show the following fact in terms of Lemmas 4.1.1 $\sim$ 4.1.3.

Lemma 4.1.4. *Let K be a holomorphic Killing field on V given by $K_u := [v, u] - b$ ($u \in V$) for some $v, b \in V$. If $K \in \mathcal{K}_{\widehat{M}^c}^h$, then we have $v \in H^1([0, 1], \mathfrak{g}^c)$ and $b = v'$.*

Proof. First we consider the case where G^c is simple. Set $\overline{K} := K - K^{\tilde{b}}$ and $w := v - \tilde{b}$, where $\tilde{b}(t) = \int_0^t b(t) dt$. From $\overline{K} = \text{ad}(w)$, we have

$$(\rho(g)_* \overline{K})_u = (\text{Ad}(g) \overline{K})_u = \text{Ad}(g)([w, g^{-1} * u]) = [\text{Ad}(g)w, u - g * \hat{0}] \quad (u \in V).$$

From this relation and (i) of Lemma 4.1.1, we have

$$\begin{aligned}
(dF_{\rho(g)*\overline{K}})_{\hat{e}}(u) &= \frac{d}{ds}|_{s=0} F_{\rho(g)*\overline{K}}(\exp su) \\
&= \frac{d}{ds}|_{s=0} \int_0^1 \text{Ad}(\exp su)((\rho(g)*\overline{K})_{\exp(-su)*\hat{0}}) dt \\
&= \int_0^1 \text{ad}(u)(\text{Ad}(g)\overline{K})_u dt \\
(4.1) \quad &= [[u, (\widetilde{\text{Ad}(g)\overline{K}}) \circ u]]_0^1 - \int_0^1 [u', (\widetilde{\text{Ad}(g)\overline{K}}) \circ u] dt \\
&= \int_0^1 [(\widetilde{\text{Ad}(g)\overline{K}}) \circ u, u'] dt \\
&= \int_0^1 [[\widetilde{\text{Ad}(g)w}, u] - [\widetilde{\text{Ad}(g)w}, g*\hat{0}], u'] dt
\end{aligned}$$

for $u \in V$. For simplicity, set $\eta := [\widetilde{\text{Ad}(g)w}, u] - [\widetilde{\text{Ad}(g)w}, g*\hat{0}]$. According to (ii) of Lemma 4.1.1, we have $\text{Im } F_K \subset T_e \widehat{M}^{\mathfrak{c}}$ and hence $\dim(\text{Span Im } F_K) \leq \dim T_e \widehat{M}^{\mathfrak{c}} < \dim \mathfrak{g}^{\mathfrak{c}}$. Since $F_{\overline{K}} - F_K$ is a constant map by (v) of Lemma 4.1.3, we have $\dim(\text{Span Im } F_{\overline{K}}) = \dim(\text{Span Im } F_K)$ and hence $\dim(\text{Span Im } F_{\overline{K}}) < \dim \mathfrak{g}^{\mathfrak{c}}$, that is, $\mathfrak{g}^{\mathfrak{c}} \ominus \text{Span Im } F_{\overline{K}} \neq \{0\}$. Take $X (\neq 0) \in \mathfrak{g}^{\mathfrak{c}} \ominus \text{Span Im } F_{\overline{K}}$. Take any $g \in \Omega_e(G^{\mathfrak{c}})$ and any $u \in T_{\hat{e}}(\Omega_e(G^{\mathfrak{c}}))$. By using (iii) of Lemma 4.1.3 and (4.1), we have

$$\begin{aligned}
\langle (dF_{\overline{K}})_g((dR_g)_{\hat{e}}(u)), X \rangle^A &= \langle (dF_{\rho(g)*\overline{K}})_{\hat{e}}(u), X \rangle^A \\
&= \int_0^1 \langle [\eta, u'], X \rangle^A dt = - \int_0^1 \langle u', [\eta, X] \rangle^A dt = - \langle u', [\eta, X] \rangle_0^A = 0,
\end{aligned}$$

where $\langle \rangle^A$ and $\langle \rangle_0^A$ are as in Section 2. From the arbitrariness of u , it follows that $[\eta, X]$ is horizontal with respect to $\phi^{\mathfrak{c}}$. Since $G^{\mathfrak{c}}$ has no center, there exists $Y \in \mathfrak{g}^{\mathfrak{c}}$ with $[X, Y] \neq 0$. Set $Z := [X, Y]$. Since $[\eta, X]$ is horizontal, it is a constant path. Hence it follows from $\langle \eta, Z \rangle^A = \langle [\eta, X], Y \rangle^A$ that $\langle \eta, Z \rangle^A$ is constant. By differentiating $\langle \eta, Z \rangle^A$ with respect to t , we have

$$(4.2) \quad \langle [\text{Ad}(g)w, u], Z \rangle^A = \langle [\text{Ad}(g)w, g*\hat{0}], Z \rangle^A \quad (u \in V).$$

Since $\mathfrak{g}^{\mathfrak{c}}$ has no center, there exists $W \in \mathfrak{g}^{\mathfrak{c}}$ with $[Z, W] \neq 0$. Since $G^{\mathfrak{c}}$ is simple, $\text{Ad}(G^{\mathfrak{c}})[Z, W]$ is full in $\mathfrak{g}^{\mathfrak{c}}$. Hence there exist $h_1, \dots, h_{2m} \in G^{\mathfrak{c}}$ such that $\{\text{Ad}(h_1)[Z, W], \dots, \text{Ad}(h_{2m})[Z, W]\}$ is a base of $\mathfrak{g}^{\mathfrak{c}}$ (regarded as a real Lie algebra), where $m := \dim_{\mathbb{C}} G^{\mathfrak{c}}$. For a sufficiently small $\varepsilon > 0$, we take $g_i \in H^1([0, 1], G^{\mathfrak{c}})$ with $g_i|_{(\varepsilon, 1-\varepsilon)} = h_i$ ($i = 1, \dots, 2m$). From (4.2), we have

$$\begin{aligned}
\langle w, \text{Ad}(h_i)[Z, W] \rangle^A|_{(\varepsilon, 1-\varepsilon)} &= \langle w, \text{Ad}(g_i)[Z, W] \rangle^A|_{(\varepsilon, 1-\varepsilon)} \\
&= -\langle [\text{Ad}(g_i^{-1})w, W], Z \rangle^A|_{(\varepsilon, 1-\varepsilon)} = -\langle [\text{Ad}(g_i^{-1})w, g_i^{-1}*\hat{0}], Z \rangle^A|_{(\varepsilon, 1-\varepsilon)} \\
&= \langle [\text{Ad}(h_i^{-1})(w|_{(\varepsilon, 1-\varepsilon)}), (g_i^{-1})'(g_i^{-1})_*^{-1}|_{(\varepsilon, 1-\varepsilon)}], Z \rangle^A = 0.
\end{aligned}$$

In particular, we have $\langle w, \text{Ad}(h_i)[Z, W] \rangle^A|_{(\varepsilon, 1-\varepsilon)} = 0$ ($i = 1, \dots, 2m$). From the arbitrariness of ε , it follows that $\langle w, \text{Ad}(h_i)[Z, W] \rangle^A = 0$ ($i = 1, \dots, 2m$). Hence we obtain $w = 0$, that is, $v = \tilde{b} \in H^1([0, 1], \mathfrak{g}^c)$ (hence $b = v'$).

Next we consider the case where G^c is not simple. Let $G^c = G_1^c \times \dots \times G_k^c$ be the irreducible decomposition of G^c and $\mathfrak{g}_i^c$ be the Lie algebra of G_i^c ($i = 1, \dots, k$). Let $\mathfrak{g}_K^c$ be the maximal ideal of $\mathfrak{g}^c$ such that the orthogonal projection of $w = v - \tilde{b}$ onto the ideal is a constant path, where we note that any ideal of $\mathfrak{g}^c$ is equal to the direct sum of some $\mathfrak{g}_i^c$'s and hence it is a non-degenerate subspace with respect to $\langle \cdot, \cdot \rangle^A$. Let $\mathfrak{g}_{\overline{K}}^c$ be the ideal corresponding to $\mathfrak{g}_K^c$ defined for $\overline{K}$. Since $\overline{K}_u = [v - \tilde{b}, u]$, we have $\mathfrak{g}_{\overline{K}}^c = \mathfrak{g}_K^c$. Now we shall show

$$(4.3) \quad (\mathfrak{g}_K^c)^\perp \subset T_e \widehat{M}^c,$$

where $(\mathfrak{g}_K^c)^\perp$ is the orthogonal complement of $\mathfrak{g}_K^c$ in $\mathfrak{g}^c$ with respect to $\langle \cdot, \cdot \rangle^A$. Let $V_i := H^0([0, 1], \mathfrak{g}_i^c)$ ($i = 1, \dots, k$). It is clear that $V = V_1 \oplus \dots \oplus V_k$ (orthogonal direct sum). The Killing field $\overline{K}$ is decomposed into $\overline{K} = \overline{K}_1 + \dots + \overline{K}_k$, where $\overline{K}_i$ is a Killing field on V_i ($i = 1, \dots, k$). For $g \in \Omega_e(G_i^c)$, we have

$$F_{\overline{K}_i}(g) = \int_0^1 \text{Ad}(g)((\overline{K}_i)_{g^{-1}*\hat{0}})dt = \int_0^1 \text{Ad}(g)(\overline{K}_{g^{-1}*\hat{0}})dt = F_{\overline{K}}(g)$$

by (i) of Lemma 4.1.1 and $\text{Ad}(g)\overline{K}_j = 0$ ($j \neq i$). Thus we have $\text{Im } F_{\overline{K}_i} = F_{\overline{K}}(\Omega_e(G_i^c)) \subset \mathfrak{g}_i^c$. From this relation and (i) of Lemma 4.1.3, we have

$$\text{Span Im } F_{\overline{K}} = \text{Span Im}(F_{\overline{K}_1} + \dots + F_{\overline{K}_k}) = \bigoplus_{i=1}^k \text{Span Im } F_{\overline{K}_i}.$$

Let $v = \sum_{i=1}^k v_i$ and $\tilde{b} = \sum_{i=1}^k \tilde{b}_i$, where $v_i, \tilde{b}_i \in V_i$ ($i = 1, \dots, k$). Since $F_{\overline{K}_i}(g) = \int_0^1 \text{Ad}(g)[v_i - \tilde{b}_i, g^{-1}*\hat{0}]dt$ ($g \in \Omega_e(G_i^c)$), $\text{Im } F_{\overline{K}_i} = 0$ when $v_i - \tilde{b}_i = 0$ and $\text{Im } F_{\overline{K}_i} = \mathfrak{g}_i^c$ when $v_i - \tilde{b}_i \neq 0$. Therefore we have $\text{Span Im } F_{\overline{K}} = \bigoplus_{i \in \{1, \dots, k\} \text{ s.t. } v_i - \tilde{b}_i \neq 0} \mathfrak{g}_i^c$. On the other hand, since $\mathfrak{g}_i^c$ is

irreducible, $(\mathfrak{g}_i^c)_{\overline{K}_i} = \mathfrak{g}_i^c$ when $v_i - \tilde{b}_i = 0$ and $(\mathfrak{g}_i^c)_{\overline{K}_i} = \{0\}$ when $v_i - \tilde{b}_i \neq 0$. Hence we have $\mathfrak{g}_{\overline{K}}^c = \bigoplus_{i \in \{1, \dots, k\} \text{ s.t. } v_i - \tilde{b}_i = 0} \mathfrak{g}_i^c$. Therefore we obtain $\text{Span Im } F_{\overline{K}} = (\mathfrak{g}_{\overline{K}}^c)^\perp$. Since $F_{\overline{K}} - F_K$ is a

constant map by (v) of Lemma 4.1.3 and $0 \in \text{Im } F_{\overline{K}}$, we have $\text{Span Im } F_{\overline{K}} \subset \text{Span Im } F_K$. Hence $(\mathfrak{g}_K^c)^\perp = (\mathfrak{g}_{\overline{K}}^c)^\perp \subset T_e \widehat{M}^c$ follows from (ii) of Lemma 4.1.1. Thus (4.3) is shown.

Next we shall show that $(R_g)_*((\mathfrak{g}_K^c)^\perp) \subset T_g \widehat{M}^c$ for any $g \in \widehat{M}^c$. Fix $g \in \widehat{M}^c$. Define $\hat{g} \in H^1([0, 1], G^c)$ with $\hat{g}(0) = e$ and $\hat{g}(1) = g$ by $\hat{g}(t) := \exp_{G^c} tX$ for some $X \in \mathfrak{g}^c$. Since $\phi^c \circ \rho(\hat{g}) = R_g^{-1} \circ \phi^c$, we have $(\phi^c)^{-1}(R_g^{-1}(\widehat{M}^c)) = \rho(\hat{g})(\widehat{M}^c)$. Since $\rho(\hat{g})_* K$ is tangent to

$\rho(\widehat{g})(\widetilde{M^c})$, in similar to (4.3), we have

$$(4.4) \quad (\mathfrak{g}_{\rho(\widehat{g})_*K}^c)^\perp \subset T_e R_g^{-1}(\widehat{M^c}) = (R_g)_*^{-1}(T_g \widehat{M^c}).$$

We have

$$(4.5) \quad (\rho(\widehat{g})_*K)_u = (\text{Ad}(\widehat{g})K)_u = [\text{Ad}(\widehat{g})v, u] - [\text{Ad}(\widehat{g})v, \widehat{g} * \hat{0}] - \text{Ad}(\widehat{g})b.$$

Denote by $\text{pr}_{\mathfrak{g}_K^c}$ the orthogonal projection of $\mathfrak{g}^c$ onto $\mathfrak{g}_K^c$. By noticing that $\text{Ad}(\widehat{g})$ preserves each $\mathfrak{g}_i^c$ (hence $\mathfrak{g}_K^c$) invariantly and that $\widehat{g} * \hat{0} = -X = -\text{Ad}(\widehat{g})X$, we have

$$\begin{aligned} & \frac{d}{dt} \text{pr}_{\mathfrak{g}_K^c} \left(\widetilde{\text{Ad}(\widehat{g})v - [\text{Ad}(\widehat{g})v, \widehat{g} * \hat{0}] - \text{Ad}(\widehat{g})b} \right) \\ &= \frac{d}{dt} \text{pr}_{\mathfrak{g}_K^c} \left(\text{Ad}(\widehat{g})(v - \widetilde{b}) + \text{Ad}(\widehat{g})\widetilde{b} - [\text{Ad}(\widehat{g})v, \widehat{g} * \hat{0}] - \widetilde{\text{Ad}(\widehat{g})b} \right) \\ &= \text{Ad}(\widehat{g})[X, \text{pr}_{\mathfrak{g}_K^c}(v - \widetilde{b})] + \text{Ad}(\widehat{g})[X, \text{pr}_{\mathfrak{g}_K^c}(\widetilde{b})] \\ & \quad + \text{Ad}(\widehat{g})\text{pr}_{\mathfrak{g}_K^c}(b) + \text{pr}_{\mathfrak{g}_K^c}[\text{Ad}(\widehat{g})v, X] - \text{Ad}(\widehat{g})\text{pr}_{\mathfrak{g}_K^c}(b) \\ &= (\text{pr}_{\mathfrak{g}_K^c} \circ \text{Ad}(\widehat{g})) \left([X, v - \widetilde{b}] + [X, \widetilde{b}] + [v, X] \right) = 0. \end{aligned}$$

Thus $\text{pr}_{\mathfrak{g}_K^c}(\text{Ad}(\widehat{g})v - [\text{Ad}(\widehat{g})v, \widehat{g} * \hat{0}] - \widetilde{\text{Ad}(\widehat{g})b})$ is a constant path. This fact together with (4.5) implies $\mathfrak{g}_K^c \subset \mathfrak{g}_{\rho(\widehat{g})_*K}^c$. By exchanging the roles of K and $(\widehat{g} * \cdot)_*K$, we have $\mathfrak{g}_{\rho(\widehat{g})_*K}^c \subset \mathfrak{g}_K^c$. Thus we obtain $\mathfrak{g}_K^c = \mathfrak{g}_{\rho(\widehat{g})_*K}^c$. Therefore $(R_g)_*(\mathfrak{g}_K^c)^\perp \subset T_g \widehat{M^c}$ follows from (4.4). Since $(R_g)_*((\mathfrak{g}_K^c)^\perp) \subset T_g \widehat{M^c}$ for any $g \in \widehat{M^c}$ and $\mathfrak{g}_K^c$ is an ideal of $\mathfrak{g}^c$, we have $\widehat{M^c} \equiv \widehat{M^c}' \times G_K^{\perp c} \subset G_K^c \times G_K^{\perp c} (= G^c)$, where $G_K^c := \exp_{G^c}(\mathfrak{g}_K^c)$ and $G_K^{\perp c} := \exp_{G^c}((\mathfrak{g}_K^c)^\perp)$. Since $\widehat{M^c}$ is irreducible and $\dim \widehat{M^c} < \dim G^c$, we have $(\mathfrak{g}_K^c)^\perp = \{0\}$, that is, $\mathfrak{g}_K^c = \mathfrak{g}^c$. This implies that $v - \widetilde{b}$ is a constant path. Therefore we obtain $b = v'$. q.e.d.

Let $\mathfrak{o}_{AK}(V)$ be the Lie algebra of all continuous skew-symmetric complex linear transformations of $(V, \langle \cdot, \cdot \rangle_0^A, \widetilde{J})$. Let $\mathfrak{g} = \mathfrak{f} + \mathfrak{p}$ be the Cartan decomposition, $\mathfrak{g}_-^c := \mathfrak{f} + \sqrt{-1}\mathfrak{p}$, $\mathfrak{g}_+^c := \sqrt{-1}\mathfrak{f} + \mathfrak{p}$ and $H_\pm^{0,c} := H^0([0, 1], \mathfrak{g}_\pm^c)$. Give $\mathfrak{o}_{AK}(V)$ the operator norm topology with respect to $\langle \cdot, \cdot \rangle_{0, H_\pm^{0,c}}^A$. Any holomorphic Killing field K on V is described as $K_u = Au + b$ for some $A \in \mathfrak{o}_{AK}(V)$ and some $b \in V$. Thus $\mathcal{K}^h$ is identified with $\mathfrak{o}_{AK}(V) \times V$. Give $\mathcal{K}^h = \mathfrak{o}_{AK}(V) \times V$ the product topology of the above topology on $\mathfrak{o}_{AK}(V)$ and the original topology of V .

Lemma 4.1.5. *The set $\mathcal{K}_{\widehat{M^c}}^h$ is closed in $\mathcal{K}^h$.*

Proof. Denote by $\overline{\mathcal{K}_{\widehat{M^c}}^h}$ the closure of $\mathcal{K}_{\widehat{M^c}}^h$ in $\mathcal{K}^h$. Take $K \in \overline{\mathcal{K}_{\widehat{M^c}}^h}$. Then there exists a sequence $\{K_n\}_{n=1}^\infty$ in $\mathcal{K}_{\widehat{M^c}}^h$ with $\lim_{n \rightarrow \infty} K_n = K$. Let $(K_n)_u = A_n u + b_n$ and $K_u = Au + b$,

where $u \in V$. From $\lim_{n \rightarrow \infty} K_n = K$, we have $\lim_{n \rightarrow \infty} A_n = A$ and hence $\lim_{n \rightarrow \infty} A_n u = Au$ ($u \in V$). Also, $\lim_{n \rightarrow \infty} b_n = b$ holds. Hence we have $\lim_{n \rightarrow \infty} (K_n)_u = K_u$ ($u \in V$). For each $u \in \widetilde{M}^c$, denote by $\text{pr}_u^\perp$ the orthogonal projection of V onto $T_u^\perp \widetilde{M}^c$. Since $\dim T_u^\perp \widetilde{M}^c < \infty$, $\text{pr}_u^\perp$ is a compact operator. Hence, since $\text{pr}_u^\perp((K_n)_u) = 0$ for all n , we obtain $\text{pr}_u^\perp(K_u) = 0$. From the arbitrariness of u , $K \in \mathcal{K}_{\widetilde{M}^c}^h$ follows. Therefore we obtain $\overline{\mathcal{K}_{\widetilde{M}^c}^h} = \mathcal{K}_{\widetilde{M}^c}^h$. q.e.d.

Take $v \in V$ and set $\widetilde{v}(t) := \int_0^t v(t) dt$. Define $g_n \in H^1([0, 1], G^c)$ ($n \in \mathbf{N}$) by $g_n(t) := \exp_{G^c}(n\widetilde{v}(t))$ and $K_n^v \in \mathcal{K}^h$ ($n \in \mathbf{N}$) by $K_n^v := \frac{1}{n}\rho(g_n)_*K$. Let $K_u = Au + b$ and $(K_n^v)_u = A_n^v u + b_n^v$. Then we have

$$\begin{aligned} (K_n^v)_u &= \frac{1}{n} \text{Ad}(g_n)(K_{g_n^{-1}*u}) = \frac{1}{n} \text{Ad}(g_n)(A(g_n^{-1} * u) + b) \\ &= \frac{1}{n} \text{Ad}(g_n) \circ A \circ \text{Ad}(g_n^{-1})u + \frac{1}{n} \text{Ad}(g_n)(A(g_n^{-1} * \hat{0}) + b) \end{aligned}$$

and hence

$$(4.6) \quad A_n^v = \frac{1}{n} \text{Ad}(g_n) \circ A \circ \text{Ad}(g_n^{-1}) \quad \text{and} \quad b_n^v = \frac{1}{n} \text{Ad}(g_n)(A(g_n^{-1} * \hat{0}) + b).$$

For $\{K_n^v\}_{n=1}^\infty$, we have the following fact.

Lemma 4.1.6. *If $K \in \mathcal{K}_{\widetilde{M}^c}^h$ and $v \in H_-^{0,c}$ with $\int_0^1 v(t) dt = 0$, then there exists a subsequence of $\{K_n^v\}_{n=1}^\infty$ converging to the zero vector field.*

Proof. Take $u \in V$. Let $u = u_- + u_+$ ($u_- \in \mathfrak{g}_-$, $u_+ \in \mathfrak{g}_+$). Then we have

$$(\text{Ad}(g_n)u_\pm)(t) = \text{Ad}(\exp_{G^c}(n\widetilde{v}(t)))u_\pm(t) = \exp(\text{ad}(n\widetilde{v}(t)))u_\pm(t) \in \mathfrak{g}_\pm^c$$

because $\widetilde{v}(t) \in \mathfrak{g}_-^c$ by the assumption. Hence we have

$$\begin{aligned} (4.7) \quad \langle \text{Ad}(g_n)u, \text{Ad}(g_n)u \rangle_{0, H_\pm^{0,c}}^A &= -\langle \text{Ad}(g_n)u_-, \text{Ad}(g_n)u_- \rangle_0^A + \langle \text{Ad}(g_n)u_+, \text{Ad}(g_n)u_+ \rangle_0^A \\ &= -\langle u_-, u_- \rangle_0^A + \langle u_+, u_+ \rangle_0^A = \langle u, u \rangle_{0, H_\pm^{0,c}}^A. \end{aligned}$$

Therefore we have $\|A_n^v\|_{\text{op}} = \frac{1}{n}\|A\|_{\text{op}} \rightarrow 0$ ($n \rightarrow \infty$) and

$$\begin{aligned} \|b_n^v\|_{0, H_\pm^{0,c}} &\leq \frac{1}{n} \left(\|A(g_n^{-1} * \hat{0})\|_{0, H_\pm^{0,c}} + \|b\|_{0, H_\pm^{0,c}} \right) \\ &= \|A\widetilde{v}\|_{0, H_\pm^{0,c}} + \frac{1}{n}\|b\|_{0, H_\pm^{0,c}} \rightarrow \|A\widetilde{v}\|_{0, H_\pm^{0,c}} \quad (n \rightarrow \infty), \end{aligned}$$

where $\|\cdot\|_{\text{op}}$ is the operator norm of $\mathfrak{o}(V)$ with respect to $\langle \cdot, \cdot \rangle_{0, H_\pm^{0,c}}^A$ and $\|\cdot\|_{0, H_\pm^{0,c}}$ is the norm of V associated with $\langle \cdot, \cdot \rangle_{0, H_\pm^{0,c}}^A$. Since $\{K_n^v \mid n \in \mathbf{N}\}$ is bounded, there exists a

convergent subsequence $\{K_{n_j}^v\}_{j=1}^\infty$ of $\{K_n^v\}_{n=1}^\infty$. Set $K_0 := \lim_{j \rightarrow \infty} K_{n_j}^v$. From $\lim_{n \rightarrow \infty} A_n^v = 0$, K_0 is a parallel Killing field on V . From $\int_0^1 v(t)dt = 0$, we have $g_n \in \Omega_e(G^\mathbf{c})$ and hence $\rho(g_n)(\widetilde{M}^\mathbf{c}) = \widetilde{M}^\mathbf{c}$. This fact together with $K \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$ deduces $K_n^v \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$. Hence have $K_0 \in \overline{\mathcal{K}_{\widetilde{M}^\mathbf{c}}^h}$. According to Lemma 4.1.5, we have $\overline{\mathcal{K}_{\widetilde{M}^\mathbf{c}}^h} = \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$. Therefore we have $K_0 \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$. Thus, since K_0 is parallel and $K_0 \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$, it follows from Lemma 4.1.4 that $K_0 = 0$. This completes the proof. q.e.d.

On the other hand, we have the following fact.

Lemma 4.1.7. *Let $K \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$ and $f \in H^0([0, 1], \mathbf{C})(= H^0([0, 1], \mathbf{R}^2))$ with $f(0) = f(1)$ and $X \in \mathfrak{g}^\mathbf{c}$. Then we have $A(fX) = [X, v]$ for some $v \in V$, where A is the linear part of K .*

Proof. Since $(\mathfrak{g}_-^\mathbf{c})^\mathbf{c} = \mathfrak{g}^\mathbf{c}$, we suffice to show in the case where $X \in \mathfrak{g}_-^\mathbf{c}$. First we consider the case of $f \in H^1([0, 1], \mathbf{C})$. Denote by f' the weakly derivative of f . Let $A(fX)(t) = u_1(t) + u_2(t)$ ($u_1(t) \in \text{Ker ad}(X)$ and $u_2(t) \in \text{Im ad}(X)$), and $u_i(t) = u_i^-(t) + u_i^+(t)$ ($u_i^-(t) \in \mathfrak{g}_-^\mathbf{c}$, $u_i^+(t) \in \mathfrak{g}_+^\mathbf{c}$) ($i = 1, 2$) and $b(t) = b^-(t) + b^+(t)$ ($b^-(t) \in \mathfrak{g}_-^\mathbf{c}$, $b^+(t) \in \mathfrak{g}_+^\mathbf{c}$). Let $g_n(t) := \exp_{G^\mathbf{c}}(nfX)$. From (4.6) and $\text{Ad}(g_n)|_{\text{Ker ad}(X)} = \text{id}$, we have

$$(4.8) \quad \begin{aligned} b_n^{f'X} &= \frac{1}{n} \text{Ad}(g_n)(A(g_n^{-1} * \hat{0}) + b) = \text{Ad}(g_n)(A(fX) + \frac{b}{n}) \\ &= u_1 + \text{Ad}(g_n)(u_2 + \frac{b}{n}). \end{aligned}$$

Since $\text{Ad}(g_n)$ preserves $\mathfrak{g}_\pm^\mathbf{c}$ invariantly and $\text{Ad}(g_n)|_{\text{Ker ad}(X)} = \text{id}$, we have

$$\begin{aligned} \langle b_n^{f'X}, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A &= \langle u_1, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A + \langle \text{Ad}(g_n)(u_2 + \frac{b}{n}), \text{Ad}(g_n)u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A \\ &= \langle u_1, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A + \frac{1}{n} \langle b, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A \rightarrow \langle u_1, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A \quad (n \rightarrow \infty). \end{aligned}$$

According to Lemma 4.1.6, there exists a subsequence $\{K_{n_j}^{f'X}\}_{j=1}^\infty$ of $\{K_n^{f'X}\}_{n=1}^\infty$ converging to the zero vector field because of $K \in \mathcal{K}_{\widetilde{M}^\mathbf{c}}^h$, $f'X \in H_-^{0, \mathbf{c}}$ and $\int_0^1 (f'X)(t)dt = (f(1) - f(0))X = 0$. Since $\lim_{j \rightarrow \infty} b_{n_j}^{f'X} = 0$, we have $\langle u_1, u_1 \rangle_{0, H_\pm^{0, \mathbf{c}}}^A = 0$, that is, $u_1 = 0$. Thus we see that $A(fX)(t) \in \text{Im ad}(X)$ holds for all $t \in [0, 1]$. That is, we have $A(fX) = [X, v]$ for some $v \in V$. Next we consider the case where f is a general element of V . Since $\overline{H^1([0, 1], \mathfrak{g}^\mathbf{c})} = V$, there exists a sequence $\{f_n\}_{n=1}^\infty$ in $H^1([0, 1], \mathfrak{g}^\mathbf{c})$ with $\lim_{n \rightarrow \infty} f_n = f$. Then, since A is continuous, we have $A(fX)(t) = \lim_{n \rightarrow \infty} A(f_n X)(t) \in \text{Im ad}(X)$ for all $t \in [0, 1]$. That is, we have $A(fX) = [X, v]$ for some $v \in V$. This completes the proof. q.e.d.

According to Lemma 2.10 of [Ch], we have the following fact.

Lemma 4.1.8. *If $B : \mathfrak{g}_-^{\mathbb{C}} \rightarrow \mathfrak{g}_-^{\mathbb{C}}$ is a linear map of the form $BX = [\mu(X), X]$ ($X \in \mathfrak{g}_-^{\mathbb{C}}$) for some map $\mu : \mathfrak{g}_-^{\mathbb{C}} \rightarrow \mathfrak{g}_-^{\mathbb{C}}$, then μ is a constant map.*

By using lemmas 4.1.7 and 4.1.8, we can show the following fact.

Lemma 4.1.9. *Let A, f, X and v be as in Lemma 4.1.7. Then v is independent of the choice of X .*

Proof. Denote by $v_{f,X}$ the above v . Define a linear map $B_1^t : \mathfrak{g}_-^{\mathbb{C}} \rightarrow \mathfrak{g}_-^{\mathbb{C}}$ by $B_1^t(X) := A(fX)(t)_{\mathfrak{g}_-^{\mathbb{C}}}$ ($X \in \mathfrak{g}_-^{\mathbb{C}}$) and a linear map $B_2^t : \mathfrak{g}_-^{\mathbb{C}} \rightarrow \mathfrak{g}_-^{\mathbb{C}}$ by $B_2^t(X) := \sqrt{-1}(A(fX)(t)_{\mathfrak{g}_+^{\mathbb{C}}})$ ($X \in \mathfrak{g}_-^{\mathbb{C}}$), where $(\cdot)_{\mathfrak{g}_{\pm}^{\mathbb{C}}}$ is the $\mathfrak{g}_{\pm}^{\mathbb{C}}$ -component of $(\cdot)$. Since $B_1^t(X) = [X, v_{f,X}(t)_{\mathfrak{g}_-^{\mathbb{C}}}]$ and $B_2^t(X) = [X, \sqrt{-1}(v_{f,X}(t)_{\mathfrak{g}_+^{\mathbb{C}}})]$, it follows from Lemma 4.1.8 that, for each $t \in [0, 1]$, $v_{f,X}(t)_{\mathfrak{g}_-^{\mathbb{C}}}$ and $v_{f,X}(t)_{\mathfrak{g}_+^{\mathbb{C}}}$ are independent of the choice of $X \in \mathfrak{g}_-^{\mathbb{C}}$. Hence $v_{f,X}$ is independent of the choice of $X \in \mathfrak{g}_-^{\mathbb{C}}$. Since $\mathfrak{g}^{\mathbb{C}}$ has no center, $v_{f,\sqrt{-1}X} = v_{f,X}$ for any $X \in \mathfrak{g}_-^{\mathbb{C}}$. Therefore $v_{f,X}$ is independent of the choice of $X \in \mathfrak{g}^{\mathbb{C}}$. q.e.d.

According to this lemma, for a fixed $K \in \mathcal{K}_{\widetilde{M}^{\mathbb{C}}}^h$, the above v depends on only f . So we denote this by v_f .

Lemma 4.1.10. *For any $f \in H^0([0, 1], \mathbb{C})$ with $f(0) = f(1)$, we have $v_f = f v_1$.*

Proof. Let $\langle \cdot, \cdot \rangle^{\mathbb{C}}$ be the complexification of the $\text{Ad}(G)$ -invariant non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$ of $\mathfrak{g}$ inducing the metric of G/K . Fix $\alpha \in \widetilde{\Delta}$ and $n \in \mathbb{N} \cup \{0\}$. Define $H_{\alpha} \in \mathfrak{h}$ by $\langle H_{\alpha}, \cdot \rangle^{\mathbb{C}} = \alpha(\cdot)$ and $c_{\alpha} := \frac{2n\pi\sqrt{-1}}{\alpha(H_{\alpha})}$. Define $g \in H^1([0, 1], G^{\mathbb{C}})$ by $g(t) := \exp_{G^{\mathbb{C}}}(tc_{\alpha}H_{\alpha})$ ($0 \leq t \leq 1$). It is clear that $g \in \Omega_e(G^{\mathbb{C}})$. Let $\overline{K} := \rho(g)_*^{-1}K$. Since $\rho(g)(\widetilde{M}^{\mathbb{C}}) = \widetilde{M}^{\mathbb{C}}$, we have $\overline{K} \in \mathcal{K}_{\widetilde{M}^{\mathbb{C}}}^h$. Let $\overline{K}_u = \overline{A}u + \overline{b}$. From Lemmas 4.1.7 and 4.1.9, there exists $\overline{v}_f \in V$ such that $\overline{A}(fX) = [X, \overline{v}_f]$ for each $f \in H^0([0, 1], \mathbb{C})$ with $f(0) = f(1)$ and each $X \in \mathfrak{g}^{\mathbb{C}}$, and $\overline{v}_f$ depends on only f . It is easy to show that $\overline{A} = \text{Ad}(g)^{-1} \circ A \circ \text{Ad}(g)$. Let $\mathfrak{h}$ be a Cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$ and $\mathfrak{g}^{\mathbb{C}} = \mathfrak{h} + \sum_{\alpha \in \widetilde{\Delta}} \mathfrak{g}_{\alpha}^{\mathbb{C}}$ be the root space decomposition with respect to $\mathfrak{h}$. Let $X \in \mathfrak{h}$ and $X_{\alpha} \in \mathfrak{g}_{\alpha}^{\mathbb{C}}$. Then we have

$$[\text{Ad}(g)\overline{v}_1, X] = [\text{Ad}(g)\overline{v}_1, \text{Ad}(g)X] = -\text{Ad}(g)(\overline{A}X) = -A(\text{Ad}(g)X) = -AX = [v_1, X].$$

It follows from the arbitrariness of X that $\text{Im}(\text{Ad}(g)\bar{v}_1 - v_1) \subset \mathfrak{h}$. Also we have

$$\begin{aligned} [\text{Ad}(g)\bar{v}_1, X_\alpha] &= \exp(2n\pi\sqrt{-1}t)^{-1}[\text{Ad}(g)\bar{v}_1, \text{Ad}(g)X_\alpha] \\ &= -\exp(2n\pi\sqrt{-1}t)^{-1}\text{Ad}(g)(\bar{A}X_\alpha) = -\exp(2n\pi\sqrt{-1}t)^{-1}A(\text{Ad}(g)X_\alpha) \\ &= -\exp(2n\pi\sqrt{-1}t)^{-1}A(\exp(2n\pi\sqrt{-1}t)X_\alpha) \\ &= \exp(2n\pi\sqrt{-1}t)^{-1}[v_{\exp(2n\pi\sqrt{-1}t)}, X_\alpha] \end{aligned}$$

and hence

$$[\text{Ad}(g)\bar{v}_1 - \exp(2n\pi\sqrt{-1}t)^{-1}v_{\exp(2n\pi\sqrt{-1}t)}, X_\alpha] = 0,$$

that is,

$$\text{Im}\left(\text{Ad}(g)\bar{v}_1 - \exp(2n\pi\sqrt{-1}t)^{-1}v_{\exp(2n\pi\sqrt{-1}t)}\right) \subset \mathfrak{z}_{\mathfrak{g}^c}(\mathfrak{g}_\alpha^c).$$

Therefore we have

$$\text{Im}\left(\exp(2n\pi\sqrt{-1}t)v_1 - v_{\exp(2n\pi\sqrt{-1}t)}\right) \subset \mathfrak{h} + \mathfrak{z}_{\mathfrak{g}^c}(\mathfrak{g}_\alpha^c).$$

From the arbitrariness of α , we obtain

$$\text{Im}\left(\exp(2n\pi\sqrt{-1}t)v_1 - v_{\exp(2n\pi\sqrt{-1}t)}\right) \subset \mathfrak{h} + \bigcap_{\alpha \in \Delta} \mathfrak{z}_{\mathfrak{g}^c}(\mathfrak{g}_\alpha^c) = \mathfrak{h}.$$

Take another Cartan subalgebra $\mathfrak{h}'$ of $\mathfrak{g}^c$ with $\mathfrak{h}' \cap \mathfrak{h} = \{0\}$. Similarly we can show

$$\text{Im}\left(\exp(2n\pi\sqrt{-1}t)v_1 - v_{\exp(2n\pi\sqrt{-1}t)}\right) \subset \mathfrak{h}'$$

and hence

$$v_{\exp(2n\pi\sqrt{-1}t)} = \exp(2n\pi\sqrt{-1}t)v_1.$$

Take any $f \in H^0([0, 1], \mathbf{C})$. Let $f = \sum_{n=-\infty}^{\infty} c_n \exp(2n\pi\sqrt{-1}t)$ be the Fourier's expansion of f . Then, since A is continuous and linear, we have $A(fX) = \sum_{n=-\infty}^{\infty} c_n A(\exp(2n\pi\sqrt{-1}t)X)$.

Also, we have $\text{ad}(X)(fv_1) = \sum_{n=-\infty}^{\infty} c_n \text{ad}(X)(\exp(2n\pi\sqrt{-1}t)v_1)$. Hence we obtain

$$\begin{aligned} [X, v_f] &= A(fX) = \sum_{n=-\infty}^{\infty} c_n A(\exp(2n\pi\sqrt{-1}t)X) \\ &= \sum_{n=-\infty}^{\infty} c_n [X, v_{\exp(2n\pi\sqrt{-1}t)}] = [X, fv_1]. \end{aligned}$$

Since $\mathfrak{g}^c$ has no center, it follows from the arbitrariness of X that $v_f = fv_1$. q.e.d.

Lemma 4.1.11. *Let K be a holomorphic Killing field on V given by $K_u = Au + b$ ($u \in V$) for some $A \in \mathfrak{o}_{AK}(V)$ and $b \in V$. If $K \in \mathcal{K}_{\widetilde{M}^c}^h$, then we have $A = \text{ad}(v)$ for some $v \in V$.*

Proof. According to Lemmas 4.1.7 and 4.1.10, there exists $v \in V$ such that $A(fX) = [v, fX]$ for any $f \in H^0([0, 1], \mathbf{C})$ with $f(0) = f(1)$ and any $X \in \mathfrak{g}^c$. Take any $u \in V$. Let $u = \sum_{n=-\infty}^{\infty} e_n \exp(2n\pi\sqrt{-1}t)$ be the Fourier's expansion of u . Then, since A is continuous and linear, we have

$$\begin{aligned} Au &= \sum_{n=-\infty}^{\infty} A(e_n \exp(2n\pi\sqrt{-1}t)) \\ &= \sum_{n=-\infty}^{\infty} [v, e_n \exp(2n\pi\sqrt{-1}t)] = [v, u]. \end{aligned}$$

Thus we obtain $A = \text{ad}(v)$.

q.e.d.

By using Lemmas 4.1.4 and 4.1.11, we shall prove Proposition 4.1.

Proof of Proposition 4.1. Let H be as in the statement of Proposition 4.1. We shall compare the Lie algebra (which is denoted by $\text{Lie } H$) of H with the Lie algebra $\rho_*(H^0([0, 1], \mathfrak{g}^c))$ of $\rho(H^1([0, 1], G^c))$. Take any $K \in \text{Lie } H(\subset \mathcal{K}^h)$. Let $\{h_s\}$ be a one-parameter transformation group of H generating K . Since h_s preserves $\widetilde{M}^c$ invariantly, K is tangent to $\widetilde{M}^c$ along $\widetilde{M}^c$, that is, $K \in \mathcal{K}_{\widetilde{M}^c}^h$. Hence, it follows from Lemmas 4.1.4 and 4.1.11 that $K = K^v$ for some $v \in V$. Thus we have $K \in \rho_*(H^0([0, 1], \mathfrak{g}^c))$. Therefore $\text{Lie } H \subset \rho_*(H^0([0, 1], \mathfrak{g}^c))$ follows. That is, $H \subset \rho(H^1([0, 1], G^c))$ follows. q.e.d.

By using Proposition 4.1, we shall prove Theorem A.

Proof of Theorem A. Without loss of generality, we may assume $\hat{0} \in \widetilde{M}^c$. Let M be as in the statement of Theorem A. According to Propositions 3.7 and 4.1, we have $\widetilde{M}^c = \rho(H) \cdot \hat{0}$ for some (connected) subgroup H of $H^1([0, 1], G^c)$. Let $\overline{H}$ be a closed connected subgroup of the anti-Kaehlerian transformation group $I_h(G^c)$ generated by $\{L_{h(0)} \circ R_{h(1)}^{-1} \mid h \in H\}$. Since $\phi^c \circ \rho(h) = (L_{h(0)} \circ R_{h(1)}^{-1}) \circ \phi^c$ for each $h \in H$, we have $\widehat{M}^c = \overline{H} \cdot e$. Set $\widehat{M} := \pi^{-1}(M)$, where π is the natural projection of G onto G/K and e is the identity element of G^c . Since $\widehat{M}$ is a component of $\widehat{M}^c \cap ((G \times G) \cdot ((e, e)\Delta G^c))$ containing e and $(\overline{H} \cap (G \times G)) \cdot ((e, e)\Delta G^c)$ is a complete open submanifold of $\widehat{M}^c \cap ((G \times G) \cdot ((e, e)\Delta G^c))$, $\widehat{M}$ is a component of $(\overline{H} \cap (G \times G)) \cdot ((e, e)\Delta G^c)$, where we note that $G^c = (G^c \times G^c)/\Delta G^c$ (hence $e = (e, e)\Delta G^c$) and that $(G \times G) \cdot ((e, e)\Delta G^c) = G(\subset G^c)$. Therefore we have $\widehat{M} = (\overline{H} \cap (G \times G))_0 \cdot ((e, e)\Delta G^c)$, where $(\overline{H} \cap (G \times G))_0$ is the identity component of

$\overline{H} \cap (G \times G)$. Set $\overline{H}_{\mathbf{R}} := (\overline{H} \cap (G \times G))_0$. Since $\widehat{M}$ consists of fibres of π , we have $\langle \overline{H}_{\mathbf{R}} \cup (e \times K) \rangle \cdot ((e, e) \triangle G^c) = \widehat{M}$, where $\langle \overline{H}_{\mathbf{R}} \cup (e \times K) \rangle$ is the group generated by $\overline{H}_{\mathbf{R}} \cup (e \times K)$. Denote by the same symbol $\overline{H}_{\mathbf{R}}$ the group $\langle \overline{H}_{\mathbf{R}} \cup (e \times K) \rangle$ newly. Set $(\overline{H}_{\mathbf{R}})_i := \{g_i \in G \mid (g_1, g_2) \in \overline{H}_{\mathbf{R}}\}$ ($i = 1, 2$), $(\overline{H}_{\mathbf{R}})'_1 := \{g \in G \mid (g, e) \in \overline{H}_{\mathbf{R}}\}$ and $(\overline{H}_{\mathbf{R}})'_2 := \{g \in G \mid (e, g) \in \overline{H}_{\mathbf{R}}\}$. It is clear that $(\overline{H}_{\mathbf{R}})'_i$ is a normal subgroup of $(\overline{H}_{\mathbf{R}})_i$. From $e \times K \subset \overline{H}_{\mathbf{R}}$, we have $K \subset (\overline{H}_{\mathbf{R}})'_2$. Since $K \subset (\overline{H}_{\mathbf{R}})'_2 \subset (\overline{H}_{\mathbf{R}})_2 \subset G$ and K is a maximal connected subgroup of G , we have $(\overline{H}_{\mathbf{R}})_2 = K$ or G and $(\overline{H}_{\mathbf{R}})'_2 = K$ or G . Suppose that $(\overline{H}_{\mathbf{R}})'_2 = G$. Then we have $\widehat{M} = G$ and hence $M = G/K$. Thus a contradiction arises. Hence we have $(\overline{H}_{\mathbf{R}})'_2 = K$. Since K is not a normal subgroup of G and it is a normal subgroup of $(\overline{H}_{\mathbf{R}})_2$, we have $(\overline{H}_{\mathbf{R}})_2 \neq G$. Therefore we have $(\overline{H}_{\mathbf{R}})_2 = K$ and hence $\overline{H}_{\mathbf{R}} \subset G \times K$. Set $\overline{\overline{H}}_{\mathbf{R}} := \{g \in G \mid (\{g\} \times K) \cap \overline{H}_{\mathbf{R}} \neq \emptyset\}$. Then, since $\widehat{M} = \overline{H}_{\mathbf{R}} \cdot ((e, e) \triangle G^c)$ and $M = \pi(\widehat{M})$, we have $M = \overline{\overline{H}}_{\mathbf{R}}(eK)$. Thus M is extrinsically homogeneous. q.e.d.

Next we prove Corollary B in terms of Theorem A.

Proof of Corollary B. Let M be as in the statement of Corollary B. According to Theorem A, M is extrinsically homogeneous. Hence it follows from Theorem A of [Koi5] that it occurs as a principal orbit of a complex hyperpolar action. Furthermore, if it admits a totally geodesic focal submanifold, then it follows from Corollary D (anf Remark 1.1) of [Koi5] that it occurs as a principal orbit of a Hermann type action. q.e.d.

References

- [BH] R. A. Blumenthal and J. J. Hebda, Complementary distributions which preserves the leaf geometry and applications to totally geodesic foliations, *Quat. J. Math.* **35** (1984) 383-392.
- [BT1] J. Berndt and H. Tamaru, Homogeneous codimension one foliations on noncompact symmetric space, *J. Differential Geom.* **63** (2003), 1-40.
- [BT2] J. Berndt and H. Tamaru, Cohomogeneity one actions on noncompact symmetric spaces with a totally geodesic singular orbit, *Tohoku Math. J. (2)* **56** (2004), 163-177.
- [BV] J. Berndt and L. Vanhecke, Curvature-adapted submanifolds, *Nihonkai Math. J.* **3** (1992), 177-185.
- [Ca] E. Cartan, Familles de surfaces isoparamétriques dans les espaces á courbure constante, *Ann. Mat. Pura Appl.* **17** (1938), 177-191.
- [Ch] U. Christ, Homogeneity of equifocal submanifolds, *J. Differential Geom.* **62** (2002), 1-15.
- [Co] H. S. M. Coxeter, Discrete groups generated by reflections, *Ann. of Math. (2)* **35** (1934), 588-621.

- [E] H. Ewert, Equifocal submanifolds in Riemannian symmetric spaces, Doctoral thesis.
- [G1] L. Geatti, Invariant domains in the complexification of a noncompact Riemannian symmetric space, *J. Algebra* **251** (2002), 619–685.
- [G2] L. Geatti, Complex extensions of semisimple symmetric spaces, *manuscripta math.* **120** (2006) 1–25.
- [HL1] E. Heintze and X. Liu, A splitting theorem for isoparametric submanifolds in Hilbert space, *J. Differential Geom.* **45** (1997), 319–335.
- [HL2] E. Heintze and X. Liu, Homogeneity of infinite dimensional isoparametric submanifolds, *Ann. of Math.* **149** (1999), 149–181.
- [HLO] E. Heintze, X. Liu and C. Olmos, Isoparametric submanifolds and a Chevalley type restriction theorem, *Integrable systems, geometry, and topology*, 151–190, AMS/IP Stud. Adv. Math. 36, Amer. Math. Soc., Providence, RI, 2006.
- [HOT] E. Heintze, C. Olmos and G. Thorbergsson, Submanifolds with constant principal curvatures and normal holonomy groups, *Int. J. Math.* **2** (1991), 167–175.
- [HPTT] E. Heintze, R. S. Palais, C. L. Terng and G. Thorbergsson, Hyperpolar actions on symmetric spaces, *Geometry, topology and physics*, 214–245 Conf. Proc. Lecture Notes Geom. Topology **4**, Internat. Press, Cambridge, Mass., 1995.
- [He] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Pure Appl. Math. 80, Academic Press, New York, 1978.
- [Hu] M. C. Hughes, Complex reflection groups, *Comm. Algebra* **18** (1990), 3999–4029.
- [KN] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Interscience Tracts in Pure and Applied Mathematics 15, Vol. II, New York, 1969.
- [Koi1] N. Koike, Submanifold geometries in a symmetric space of non-compact type and a pseudo-Hilbert space, *Kyushu J. Math.* **58** (2004), 167–202.
- [Koi2] N. Koike, Complex equifocal submanifolds and infinite dimensional anti-Kaehlerian isoparametric submanifolds, *Tokyo J. Math.* **28** (2005), 201–247.
- [Koi3] N. Koike, Actions of Hermann type and proper complex equifocal submanifolds, *Osaka J. Math.* **42** (2005) 599–611.
- [Koi4] N. Koike, A splitting theorem for proper complex equifocal submanifolds, *Tohoku Math. J.* **58** (2006) 393–417.
- [Koi5] N. Koike, Complex hyperpolar actions with a totally geodesic orbit, *Osaka J. Math.* **44** (2007), 491–503.
- [Koi6] N. Koike, The homogeneous slice theorem for the complete complexification of a proper complex equifocal submanifold, *arXiv:math.DG/0807.1595v1*.
- [Koi7] N. Koike, The complexifications of pseudo-Riemannian manifolds and anti-Kaehler geometry, *arXiv:math.DG/0807.1601v1*.
- [Koi8] N. Koike, On curvature-adapted and proper complex equifocal submanifolds, *arXiv:math.DG/0809.4933v1*.
- [Kol] A. Kollross, A classification of hyperpolar and cohomogeneity one actions, *Trans. Amer. Math. Soc.* **354** (2002), 571–612.
- [O’N] B. O’Neill, *Semi-Riemannian Geometry, with applications to relativity*, Pure Appl. Math. 103, Academic Press, New York, 1983.
- [OS] T. Ohshima and J. Sekiguchi, The restricted root system of a semisimple symmetric pair, *Group representations and systems of differential equations* (Tokyo, 1982), 433–497, Adv. Stud. Pure Math. **4**, North-Holland, Amsterdam, 1984.

- [P] R. S. Palais, Morse theory on Hilbert manifolds, *Topology* **2** (1963), 299–340.
- [PT1] R. S. Palais and C. L. Terng, A general theory of canonical forms, *Trans. Amer. Math. Soc.* **300** (1987), 771–789.
- [PT2] R. S. Palais and C. L. Terng, Critical point theory and submanifold geometry, *Lecture Notes in Math.* **1353**, Springer-Verlag, Berlin, 1988.
- [S1] R. Szöke, Involutive structures on the tangent bundle of symmetric spaces, *Math. Ann.* **319** (2001), 319–348.
- [S2] R. Szöke, Canonical complex structures associated to connections and complexifications of Lie groups, *Math. Ann.* **329** (2004), 553–591.
- [Te1] C. L. Terng, Isoparametric submanifolds and their Coxeter groups, *J. Differential Geom.* **21** (1985), 79–107.
- [Te2] C. L. Terng, Proper Fredholm submanifolds of Hilbert space, *J. Differential Geom.* **29** (1989), 9–47.
- [Te3] C. L. Terng, Polar actions on Hilbert space, *J. Geom. Anal.* **5** (1995), 129–150.
- [TT] C. L. Terng and G. Thorbergsson, Submanifold geometry in symmetric spaces, *J. Differential Geom.* **42** (1995), 665–718.
- [Th] G. Thorbergsson, Isoparametric foliations and their buildings, *Ann of Math.* **133** (1991), 429–446.
- [W] B. Wu, Isoparametric submanifolds of hyperbolic spaces, *Trans. Amer. Math. Soc.* **331** (1992), 609–626.